Vol. 33 No 12

BULLETIN

OF THE

TORREY BOTANICAL CLUB

DECEMBER, 1906

American ferns — VII

LUCIEN MARCUS UNDERWOOD

A. THE AMERICAN SPECIES OF STENOCHLAENA

A short time since we presented an account of the species of *Stenochlaena* from the Old World,* stating that the American species would form the subject of a later paper. We present herewith a synopsis of the known American species, all of which belong to the section *Lomariopsis*. Instead of ten species as estimated at that time, we are forced to recognize twelve; besides these there are several collections which do not seem to fall within the limits of any of the described species, and we have set these aside for further information dependent on more extended exploration in the regions where they have already been collected a single time. A list of these possible candidates for recognition as species is given at the end of this paper under the *Species inquirendae*. All the following species are included under *Acrostichum sorbifolium* in the *Synopsis Filicum* of Hooker and Baker, as was also the case with the greater part of the species of the Old World as discussed and in part figured in our previous paper. One purpose of binomial names is to enable botanists to refer definitely to some particular well-defined plant. There are some, commencing with Goethe, who maintain that this is the only *raison d'être* of species in the text-books anyway. Now if a single name like *Acrostichum sorbifolium* is allowed to stand for a series of twenty-five distinct and recognizable groups of plants, each found in some definite quarter of the tropical world, somewhere from Florida to Brazil,

* Bull. Torrey Club 33: 35–50. 1906.

[The BULLETIN for November 1906 (33: 541–590, *pl. 18–26*) was issued 19 D 1906.]

around the world to Sierra Leone and Mauritius, thence to India, Java, the Philippines, New Zealand, and Samoa, each with distinctive characters and each with a well-defined geographic range, how is one to know how to refer *definitely* to any particular group of this series, for example the one figured in this paper (FIGURES 1–3) which has a very limited distribution in eastern Cuba, together with a very pronounced type of leaf that no one would possibly mistake for any other? Or how shall we refer to another very definite plant of the lowlands of Florida and Cuba which we have recently seen fit to describe as *Stenochlaena Kunzeana* * and which is figured in this paper (FIGURES 4–6)? It is true they could be referred to as varieties of *Stenochlaena sorbifolia;* so could our species of soft maples be referred to as varieties of the sugar maple, but I believe none of our so-called conservatives has thought of such a course, even though the foliage of the maples is really much closer than the two plants in question. Besides, we do not know which is the parent stock and which is the variation, or better we do know that both have diverged from a common ancestor in the past so that neither one could with accuracy be said to be a variety of the other. Being distinct recognizable entities, we prefer to call these units of classification *species*, the same as we do our well recognized species of maples and other plants of a similar sort.

Linnaeus founded *Acrostichum sorbifolium* on two things that turn out to be distinct. One was a plant figured by Plumier from Martinique (afterwards copied by Petiver), and the other was a plant figured by Sloane from Jamaica. The acquisition of the extensive herbarium of Père Duss from the islands of Martinique and

* We are gravely informed by the confident editor of the Fern Bulletin (14: 95) that this species is *Lomaria procera!* and that "the [Florida] plant reported by A. A. Eaton as *Acrostichum sorbifolium* is also omitted" [*i. e.*, in the last number of this series of papers]. *Lomaria procera* was originally described from New Zealand and like its congeners of the mountain regions of the West Indies is a terrestrial fern growing in a crown after the manner of our native Christmas fern, and like other species of *Lomaria* possesses a distinct linear intramarginal sorus covered by an indusium; while *Stenochlaena Kunzeana*, originally designated (but not described) by Presl seventy years ago from the lowlands of Cuba, is a climbing fern with scattered leaves, creeps up the trunks of trees three or four meters, and like other members of its genus possesses no indusium and not even a definite sorus. Mere details (?) of structure, habit, geographic and altitudinal distribution like these, seem to amount to nothing in the mind of the editor of "the only fern magazine in the world."

Guadeloupe places the New York Botanical Garden in possession of the most complete series of plants from those islands that has ever been brought together. Specimens of *Stenochlaena* in this herbarium, as well as older collections from Martinique seen in European herbaria, when compared with the recent extensive collections in the island of Jamaica, have enabled us clearly to differentiate the plant of the Lesser Antilles, which extends also into Porto Rico and Hispaniola, from the Jamaican plant, more or less common in the hill region of that island.

Each one of the species described or recognized in this paper has a very definite geographic area as determined from a study of practically all the material that has yet been collected. The species are most abundantly represented in our own collections, but the materials at Vienna, Prague, Berlin, Paris, Kew, Washington and New Haven have contributed type specimens and additional information.

The American species which we recognize may be separated by the following synopsis:

Pinnae tapering gradually in the lower third, truncate-acuminate at the apex; sporophyllary pinnae narrow (3–4 mm.). 12. *S. Wrightii.*

Pinnae blunt or cuneate at base, but never long-tapering below.

Pinnae mostly 2–4 cm. wide; sporophyllary pinnae 7–16 mm. wide.

Pinnae few (1–5). 8. *S. Prieuriana.*

Pinnae more numerous (7–20 or more).

Sporophyllary pinnae 12–16 mm. wide; scales of the rootstock and bases of the leaf-stalks brownish-black. 5. *S. japurensis.*

Sporophyllary pinnae 7–8 mm. wide.

Scales of the rootstock and bases of the leaf-stalks narrowly lanceolate, ferruginous; sporophyllary pinnae mostly 12 cm. long. 2. *S. erythrodes.*

Scales of the rootstock and bases of the leaf-stalks broad, ovate, pallid; sporophyllary pinnae shorter, 6–9 cm., stalked. 4. *S. jamaicensis.*

Pinnae normally under 2 cm. wide; sporophyllary pinnae under 5 mm. wide (except in no. 7).

Pinnae irregularly sharply serrate, not over 1 cm. wide, cuneate at the base. 6. *S. Kunzeana.*

Pinnae entire, undulate, or slightly and uniformly serrulate or denticulate.

Pinnae truncate at the base.

Pinnae 7–8 cm. or more long; sporophyllary pinnae 3 mm. wide. 3. *S. Fendleri.*

Pinnae 4–7 cm. long; sporophyllary pinnae under 2 mm wide. 11. *S. vestita.*
Pinnae cuneate at base (acute or obtuse).
Pinnae 4–8 cm. long.
Pinnae 20–25-jugate; sporophyllary pinnae nearly sessile; scales of the rootstock long-hairy ciliate. 1. *S. angusta.*
Pinnae 10–15-jugate; sporophyllary pinnae stalked; scales of the rootstock smooth. 10. *S. sorbifolia.*
Pinnae 10–15 cm. long, tapering to a long-acuminate apex.
Sporophyllary pinnae 5–6 mm. wide, 9 cm. long, long-stalked. 7. *S. Maxoni.*
Sporophyllary pinnae not over 2 mm. wide, 10–12 cm. long, short-stalked. 9. *S. recurvata.*

1. **Stenochlaena angusta** sp. nov.

Wide-climbing on trees. Roostock rather slender, 4–8 mm. thick, densely covered at the apex with pale, narrowly lanceolate, long-ciliate scales, becoming naked below; leaves scattered, narrow, tapering at both ends, 45–60 cm. long, with a short stalk (6–8 cm.) which is densely covered below with scales like those of the rootstock, and with still narrower scattered fibrils which extend up the rachises; pinnae 20–25-jugate, obtusely cuneate at base forming a very short stalk, 6–7 cm. long, 10–12 mm. wide, the margins finely and deeply undulate, tapering at the more or less falcate apex into an acuminate point; veins simple or once forked, rather close, with 11–15 intercostal spaces to 1 cm.; sporophyls slightly shorter than the leaves, the pinnae 22–25-jugate, 55–8 cm. long, 3–4 mm. wide, obtuse at the base with a short stalk, tapering gradually to the apex. [FIGURE 6.]

COLOMBIA: Jordan, Prov. Santa Marta, *H. H. Smith 1051* (type). "Creeping on tree trunks to 15 feet or more. Rare in deep forest, 1200–1400 feet."

To the same species we should refer *Funck 774* collected in 1845 (P) * and *Spruce 1556* collected at Tarapoto, Peru (K). Also more or less fragmentary recent collections from Bolivia: Lower Rio Cocos, *Williams 1190;* San Raphael, *Williams 1189;* and more doubtfully the juvenile forms, Isapuri, *Williams 1203.*

* Unless otherwise stated, specimens cited are in the collections in the New York Botanical Garden. For convenience other collections are marked with an initial: K = Kew, B = Berlin, P = Paris, Pr = Prague, V = Vienna, etc.

2. **Stenochlaena erythrodes** (Kunze).

Acrostichum erythrodes Kunze, Flora **22**[1]: Beibl. 46. 1839. (Type from Ilheos, Brazil, *Herb. Mart. 366.*)

RANGE: Brazil, Prope Ilheos, *Herb. Mart. 366*; *Sellow* (B); Organ Mountains, *Gardner* (?) *101* (K).

In *Index Filicum*, Mr. Christensen refers this plant with two other species of the genus to *Stenochlaena marginata* (Schrad.) C. Chr., the original of which was described by Schrader in 1824 in his report on the ferns collected in Brazil by Maximilian, Prince of Neuwied, as follows: "[LOMARIA] MARGINATA: *frondibus pinnatis; sterilium pinnis oblongo-acuminatis subintegerrimis basi superiori cuneatis glabris indusiis crenato-incisis rhachi marginata.*" Goett. Gel. Anz. 871. 1824. In the absence of the type and with such a description it would seem safer to leave this name in the waiting list until more certainty is attainable in regard to its status; if the description is correct, the plant is not even a *Stenochlaena*. In any case the three united species have little in common and can easily be distinguished, when real specimens are at hand.

3. **Stenochlaena Fendleri** (D. C. Eaton).

Lomariopsis Fendleri D. C. Eaton, Mem. Am. Acad. II. **8**: 195. 1860. (Type from Venezuela, *Fendler 335.*)

RANGE: Known only from its type locality. [FIGURES 9 and 10.]

4. **Stenochlaena jamaicensis** sp. nov.

Wide-climbing on tree-trunks. Rootstock 1.5–2 cm. in diameter, densely covered with pallid ciliate scales which are broadly lanceolate to lanceolate-ovate in shape, 2–3 mm. wide; leaves 50 cm. or more long on stramineous stalks 12–14 cm. long, densely covered at the base with scales similar to those of the rootstocks, simply pinnate, 5–10-jugate, the pinnae obtusely cuneate at the base, more or less abruptly tapering at the apex, 8–12 cm. long, 2–2.5 cm. wide, the margins slightly repand and very minutely revolute in drying; veins simple or occasionally forked, inconspicuous, with about 10–11 intercostal spaces to 1 cm.; sporophyls 35 cm. or more long, on similar but shorter (8–10 cm.) leaf-stalks, about 9-jugate; pinnae short-stalked (3–4 mm.), obtuse at base, short-tapering at the apex, 7–8 cm. long by 7–8 mm. wide. [FIGURES 13 and 14.]

JAMAICA: Vicinity of Mandeville, *Maxon 2566* (type, in the United States National Herbarium and the New York Botanical Garden); Manchester, *Day;* no locality, 1884–5, *Balch.*



Normal pinnae of American species of *Stenochlaena:* 1–3, *S. Wrightii.* 4, 5, *S. Kunzeana.* 6, *S. angusta.* All the figures natural size.

Besides the above, the following mostly sterile plants appear to go here: Hollymount, *Maxon 1904, 2217;* Tyre, Cockpit country, *Underwood 3305;* Mansfield, near Bath, *Underwood 2782.*

Young forms collected with the last number seem to indicate a widely different juvenile condition from that which obtains in the



Normal pinnae of American species of *Stenochlaena*: 7, 8, *S. vestita*. 9, 10, *S. Fendleri*. 11, 12, *S. Maxoni*. 13, 14, *S. jamaicensis*. All the figures natural size.

common Windward Island species (*S. sorbifolia*) from which this is a segregation. In the last-named species the juvenile form is

so unlike the mature condition as often to be regarded as belonging to an entirely different species. In the present species the juvenile leaves fail to show this strongly dimorphic character. Further material bearing on this point is a desideratum.

5. STENOCHLAENA JAPURENSIS (Mart.) Griseb. Fl. Brit. W. I. 676. 1864.

Acrostichum japurense Mart. Ic. Pl. Crypt. Bras. 86. *pl. 24.* 1834. (Type from Japura River, N. Brazil.)

Lomariopsis phlebodes Kunze, Linnaea **9**: 33. 1835. (Type from Maynas, Peru, *Poeppig.*)

RANGE: Guatemala and Costa Rica to Trinidad, Guiana, and Brazil and down the Andes as far as Bolivia and Peru. Apparently a lowland species. We have specimens as follows:

GUATEMALA: Cubilquitz (Alta Verapaz) *von Tuerckheim* (*J. D. S. 8043*).

COSTA RICA: Turrialba, *Maxon 184.*

TRINIDAD: *Fendler 106.*

PANAMA: *Hayes 4.*

GUIANA: Essequibo, *Jenman 2312;* Mt. Ragwa, *Jenman;* Demerara River, *Jenman.*

COLUMBIA: San Pablo, *Holton 21;* Santa Marta, *H. H. Smith 2697.*

BOLIVIA: Tumupara, *Williams 1192;* San Buena Ventura, *Williams 1196.*

The species does not appear to have been collected much below the Amazon in Brazil, but there are specimens at Kew collected in the Upper Amazon, *Traill 1394*, and Para, *Spruce 27* and *569.* We have seen Poeppig's plant (cotype of *Lomariopsis phlebodes*) at Vienna and it does not appear to be separable from the widely distributed species as listed above. This species and *S. erythrodes* of Southern Brazil are united under a distinct name (noted above) by Christensen in his *Index Filicum.*

6. STENOCHLAENA KUNZEANA (Presl) Underw. Bull. Torrey Club **33**: 196. 1906. (Type from Cuba, *Wright 973.*)

Olfersia Kunzeana Presl, Tent. Pterid. 235. 1836 (*nomen nudum*).

RANGE: Cuba, Hispaniola, and southern Florida. (FIGURES 4 and 5.)

Additional stations to those given with the original description are as follows:

CUBA: El Guama (Prov. Pinar del Rio), *Palmer & Riley 392;* Sevilla Estate near Santiago, *Taylor 473.*

7. **Stenochlaena Maxoni** sp. nov.

Climbing on tree-trunks. Rootstock wide-creeping, 8–10 mm. in diameter, densely covered with pale straw-colored, narrowly lanceolate scales, which are usually less than 1 mm. wide; leaves scattered, 65 cm. or more long, on long, rather stout leaf-stalks, 25 cm. long, which have scattering scales throughout like those of the rootstock intermixed with brown fibrils which extend upwards through the rachises; pinnae about 9-jugate, distant, rather long-stalked (5–7 mm.), acutely cuneate at the base, long-acuminate at the apex, 12–14 cm. long, about 2 cm. wide, the margins irregularly undulate, very slightly reflexed in drying; veins very prominent, converging in pairs at the base when simple, occasionally forked, somewhat spaced, with about 8 intercostal spaces to 1 cm.; sporophyls 55 cm. or more long, with leaf-stalks similar to those of the leaves, about 10-jugate, the pinnae distant, stalked (4–6 mm.), obtuse at base, tapering at apex, 9 cm. long, 5–6 mm. wide. [FIGURES 11 and 12.]

COSTA RICA: La Palma, alt. 1450–1550 m., *Maxon 411.*

It is with pleasure that we name this clearly marked species from Costa Rica for Mr. William Ralph Maxon through whose discriminating and careful field study we know so much of the real characters of the fern flora of Costa Rica. The practice of sending specialists into the field is of the greatest importance if we ever expect to secure accurate information regarding the real characters of plants.

8. **Stenochlaena Prieuriana** (Fée).

Lomariopsis Prieuriana Fée, Mém. Foug. **2**: 66. *pl. 25. f. 1.* 1845. (Type from Guiana, *Herb. Richard.*)

RANGE: Known only from Guiana, *Appun 128* (K); Pomeroon River, *Jenman;* Potaro River, *Jenman;* Ragwa, *Jenman.*

This species has been combined with *Stenochlaena japurensis,* with which it has closest alliance; its limited number of pinnae is apparently a constant character, and until this is found to be due to accidental causes it may best remain distinct as Fée left it.

9. **Stenochlaena recurvata** (Fée).

Lomariopsis recurvata Fée, Mém. Foug. **2**: 68. *pl. 28.* 1845. (Type from Teapa, Mexico, *Herb. Delessert et Mus. Paris.*)

RANGE: Known only from its type collection, of which we have seen the specimen preserved at Paris. A very distinct species.

10. STENOCHLAENA SORBIFOLIA (L.) J. Sm. Jour. Bot. **4**: 149. 1841. *Acrostichum sorbifolium* L. Sp. Pl. 1069. 1753. (Type from Martinique, based on Plumier *pl. 117.*)

RANGE: Lesser Antilles, Porto Rico, and Hispaniola.

We have seen specimens of this characteristic species, whose distribution has been erroneously supposed to be well-nigh universal, from the following islands:

MARTINIQUE: *Duss 1698.*

GUADELOUPE: *Duss 4139.*

DOMINICA: *Lloyd 344, 743, 744, 869.*

ST. VINCENT: *Guilding* (K), *Chickley* (B).

ST. KITTS: *Hon. J. H. H. Berkley, Britton & Cowell 454.*

PORTO RICO: 1824, *Balbis* (P), *Schwanecke* (K), *Booth* (K), *Underwood & Griggs 843, Britton & Marble 1084.*

HISPANIOLA: Petit Borgne, *Nash 468.*

11. **Stenochlaena vestita** (Fourn.). [FIGURES 7 and 8.]

Lomariopsis vestita Fourn. Bull. Bot. Soc. France **19**: 250. 1872. (Type from Chontales, Nicaragua, *Levy 476.*)

Acrostichum Pittieri Christ, Bull. Soc. Bot. Belg. **35**: 243. 1896. (Type from Costa Rica, *Pittier 6931.*)

Lomariopsis guatemalensis Presl, in herb. (Sterile plant only. "In Guatimala ad ripam occidentalem fluvii S. Juan, *Friedrichsthal.*")

Stenochlaena Pittieri Diels, Die Nat. Pflanzenf. **1**[4]: 251. 1899.

RANGE: Apparently throughout Central America. We have seen specimens as follows:

GUATEMALA: Alta Verapaz, *von Tuerckheim* (*J. D. S. 8042*), San Juan River,* *Friedrichsthal* (V, Pr).

NICARAGUA: Chontales, *Levy 496* (K, P), *Tate 72* (K, two sheets).

COSTA RICA: La Concepcion, *J. D. S. 6942; Wercklé;* Turrialba, *Maxon 163.*

* There appears to be some doubt whether this station is in Guatemala or not.

PANAMA: Lion Hill Station, *Hayes 379* (K).

This species, which was first named (though not published) by Presl sixty years or more ago, and twice published since, is rather close to *S. Fendleri*, and the Panama specimen cited above suggests a possible transition between the two. In the absence of any more definite light on this subject it is wiser to keep them separate for the present.

12. STENOCHLAENA WRIGHTII (Mett.) Griseb. Cat. Pl. Cub. 277. 1866.

Lomariopsis Wrightii Mett.; D. C. Eaton, Mem. Am. Acad. II. 8: 195. 1860. (Type from Monte Verde, eastern Cuba, *Wright 787*.)

RANGE: Confined to eastern Cuba. [FIGURES 1–3.]

This most peculiar species has been twice collected since Wright's time at a second station still farther eastward than Monte Verde, *viz.*: at Baracoa, *Pollard & Palmer 220*, and on the slopes of El Yunque, near Baracoa, *Underwood & Earle 948*.

SPECIES INQUIRENDAE

1. We have collected in Jamaica what appears to be a very distinct species with small pinnae (4–5.5 cm. by 1.2–1.5 cm.) which are about 13-jugate on short stalks, acutely tapering at the base, and the rootstocks densely covered with dark ferruginous narrowly lanceolate ciliated scales, 1 mm. wide. The veins are very close (12–15 intercostal spaces to 1 cm.). Unfortunately only immature sporophyls have been found; hence our hesitancy to describe what appears to be distinct. The plants were collected in the Cockpit country near Troy, *Underwood 2866* and *2887* (the latter sterile only).

The present genus illustrates excellently the doubt often occasioned by improper field work. There are in various collections a series of scrappy specimens often consisting of the upper half of a leaf only or sometimes with a similar upper portion of a sporophyl unconnected with the leaf, and neither showing anything of the character of the stem, which in this genus, as in most others, presents most important diagnostic specific characters. To break off the mere tip of a fern as a representative of a species is as idi-

otic a proceeding as to break off the upper half of a compound leaf of ash or hickory or sumac and expect a botanist to describe a species from this fragment only. And yet a very large number of fern species have been described from just such fragments and their types are often of such a character that after real personal field work in the type locality of a given species, it is only with the greatest uncertainty that we can recognize the elegant and complete specimens of modern collecting as the same as these old fragments, especially after they have been handled for two or three generations. We have a series of these fragments from Jamaica and elsewhere that are now in doubt and will very likely remain in this condition to the end of time. One hesitates to throw away an imperfect specimen after it has once been mounted in a collection, but it might be better except, of course, in the case of old types to commit them to the flames and save succeeding generations from wasting gray matter on them.

2. *Fendler 121*, from Trinidad (distributed by Professor D. C. Eaton as *Acrostichum sorbifolium*), probably represents a distinct species. We had long supposed that this represented *Lomaria longifolia*, a species described by Kaulfuss in 1824, citing: (1) Willdenow's description of *Acrostichum sorbifolium* (Sp. Pl. 5: 115, *excl. syn.*), and (2) a figure of a separate pinna by Plumier on plate 117 (the type of *Acrostichum sorbifolium* L.). An examination of Willdenow's herbarium shows that he had under this species a plant with elongate pinnae, which seems to differ in no other way from the ordinary *Stenochlaena sorbifolia*, and analogous specimens appear in Père Duss' collections both from Martinique and Guadeloupe, as well as in those of Sherring from Grenada, and among those of Lloyd from Dominica.

The Trinidad plants differ more widely (1) in the vestiture of the bases of the leaf-stalks and the rootstocks, the scales being densely long-hairy and ciliate, while they are smooth in *S. sorbifolia*; (2) in the form of the pinnae and their basal characters; (3) in the closeness of the venation, there being 11–12 intercostal spaces to 1 cm. (instead of 7–9 as in *S. sorbifolia*); and in various other characters. Unless further information appears, *Lomaria longifolia* Kaulf. will become a synonym of *S. sorbifolia*, but the Trinidad plants may come to be recognized as a distinct species of *Stenochlaena*.

3. A single specimen from Potrero, Mexico, *Hahn 56*, in herb. Paris, indicates a species distinct from any known. The pinnae are 9 by 2.5 cm., with prominent veins, while the sporophyllary pinnae are narrowly linear, 13 cm. long by 3 mm. wide. The specimens were distributed as *S. erythrodes* but of course have nothing to do with that South Brazilian species. Potrero is in the state of Hidalgo.

4. Two specimens in herb. Berol.: Bahia, *Luchnath 86*, and *Sellow* (without locality), were marked by Klotzsch as a new species in *Olfersia* but apparently have not been described. They represent a species of *Stenochlaena* which recalls *S. Maxoni* as described above, but have marked differences. The pinnae are acute at base but are sessile, 11–12 by 2 cm., tapering to an acuminate point; the sporophyllary pinnae are 6–7.5 cm. long by 4 mm. wide and also sessile. The two specimens are not quite counterparts of each other, the Luchnath specimen having the sporophyllary pinnae up to 9 cm. long. They probably represent a good species which had best await further exploration in the little-known regions of Brazil.

5. A second specimen from Brazil, collected by Sellow, also in herb. Berol., bears a second unpublished name by Klotzsch under *Olfersia*. It has pinnae rounded at the sessile base, 17 by 2.5 cm., tapering gradually to the apex; the sporophyllary pinnae are 10 cm. long by 6 mm. wide; the veins are close, 12 or more intercostal spaces to 1 cm. This plant is not like any described species and is probably new, as indicated in Klotzsch's herbarium name.

The genus *Olfersia*, containing the common *Olfersia cervina* of the American tropics and a second Brazilian species which is the type of the genus (*O. corcovadensis*), is characterized by its distinct marginal vein with which the other veins unite; its sporophyls are similar to those of *Polybotrya* but simpler. It is not a near ally of *Stenochlaena*.

B. The status of Poecilopteris crenata Presl

The genus *Leptochilus* was founded by Kaulfuss in 1824, based on *Acrostichum axillare* Cav. This long-known plant possesses a simple leaf with the complex venation of *Drynaria*, *Tectaria*, and *Phymatodes*. Whether all the species with this type of anas-

tomosing veins, formerly associated erroneously under the genus *Gymnopteris* (*Acrostichum* § *Gymnopteris* of *Synopsis Filicum*), will form a single genus, or be differentiated into several, as held by Presl, John Smith and Fée, may well form a later discussion. Suffice it to say that the American species, elaborately treated by Carl Christensen under *Leptochilus* § *Bolbitis*,* in our opinion are not congeneric with *Acrostichum axillare*, and as a coherent group must bear the name *Poikilopteris* † as the oldest name applied to any species included within the present generic limits. Their venation is amply characterized in the illustrations given in Mr. Christensen's admirable paper and may be seen in the accompanying figure (FIGURE 16). We do not purpose a duplication of Mr. Christensen's work, but having seen nearly all the original specimens cited in his paper, and in addition the equally extensive collection at Kew which materially supplements those cited, and particularly the originals of Presl at Prague, we wish to make a correction regarding the status of one of the doubtful species cited at the close of his paper.



Poecilopteris crenata Presl. Drawn from Presl's type material at Prague. 15, Sporophyllary pinna. 16, Normal sterile pinna. Both natural size.

Mr. Christensen says (page 297): "*Poecilopteris crenata* Presl, Epim. bot. 174. 1849.—Rio. After a remark on a sheet in Herb. Berol. (by Mettenius ?) and after the description this is only a common form of *L. serratifolius*."

The above quotation notes the usual European neglect to look up type specimens so easily accessible as those in Presl's herbarium, and also emphasizes the force of a statement that cannot be

* Bot. Tidsskrift **26**: 283–300. 1904.

† Eschweiler, Linnaea **2**: 117. 1827; afterwards (1836) called *Poecilopteris* by Presl.

reiterated too often, *viz.*: "A species is not a description, but a group of allied individuals," in this case represented by a single well-preserved type specimen. The accompanying figures (15–16) are drawn directly from Presl's *Poecilopteris crenata* and it will be seen to be, not the plant to which it has been erroneously referred by those who have never seen it, but instead a species closely allied to the one described in Mr. Christensen's paper as *Leptochilus contaminoides* (Christ) C. Chr. Having seen the type number of the Paraguay plant both at Berlin and at Kew, and the Brazilian specimens also referred by Mr. Christensen to Christ's species, we are inclined to regard the Paraguay plant as distinct from the others, and refer all the Brazilian plants cited to Presl's species. Otherwise Dr. Christ's species must fall under it in synonymy. We present the following comparisons of *Balansa 2852* and Presl's plant:

Balansa 2852 (Type of *Gymnopteris contaminoides* Christ).	*Poecilopteris crenata* Presl. (*Cf.* FIGURES 15 and 16.)
Pinnae 3.5 cm. wide.	Pinnae 2 cm. wide.
Color bright-green.	Color distinctly red.
Margins regularly crenate-toothed.	Margins irregularly undulate.
Areolae in 5 (4–6) rows between the costae.	Areolae in 4 (3–5) rows between the costae.
Sporophyllary pinnae 8 mm. wide.	Sporophyllary pinnae 6 mm. wide.

At a later time we hope to add something to the series of species so well described by Mr. Christensen, when the other species may be more appropriately transferred to what we regard their correct generic name, *Poikilopteris*.

COLUMBIA UNIVERSITY.

Some desmids from Newfoundland

Joseph Augustine Cushman

During the past year material has been obtained from Newfoundland which has added many species to the few which were previously known to belong to the flora of the island. The greatest number of species were obtained from material collected by Mr. Owen Bryant at St. Anthony in May, 1906. Some of the species were not previously known from North America. In addition to this material a number of species were obtained from *Utricularia* collected by Waghorne at Bay of Islands. This added several species not obtained in any of the other material. Included in the present paper are the species found by the writer in Dr. G. M. Allen's material from Rose au Rue, and published in this journal for November, 1904. In certain cases the synonymy has been altered and in these species the corrections are here made. Material has thus been obtained from three points on the island fairly remote from one another. As a result of the work seventy-two species included in seventeen genera are here noted from Newfoundland. None were recorded from the island previous to the former paper.

MESOTAENIUM Näg. 1849

1. Mesotaenium DeGrevi Turn.
   Length 83 μ; breadth 20 μ. Occasional, St. Anthony.
2. Mesotaenium Endlicherianum Näg.
   Length 35 μ; breadth 12 μ. Very common, Rose au Rue.

CYLINDROCYSTIS Menegh. 1838

3. Cylindrocystis Brebissonii Menegh.
   Length 54 μ; breadth 18 μ. Fairly common, St. Anthony.
4. Cylindrocystis americana minor Cushman.
   Length 40 μ; breadth 18 μ. Common, Rose au Rue.

5. CYLINDROCYSTIS DIPLOSPORA Lund.

Length 71 μ; breadth 37 μ; isthmus 33 μ. Occasional, St. Anthony.

NETRIUM Näg. 1849

6. NETRIUM DIGITUS (Ehrenb.) Itzigs. & Rothe.

Length 160–190 μ; breadth 63–80 μ. Common at Rose au Rue and St. Anthony. The prevailing form is short and much swollen at the central portion, thence narrowing fairly rapidly to the apex.

7. NETRIUM NAEGELII (Bréb.) W. & G. S. West.

Length 140–145 μ; breadth 37 μ. Occasional, Bay of Islands and St. Anthony.

PENIUM Bréb. 1844

8. PENIUM LIBELLULA INTERMEDIUM Roy & Biss.

Length 98–110 μ; breadth 24–29 μ. Occasional, Rose au Rue.

CLOSTERIUM Nitzsch. 1817

9. CLOSTERIUM CYNTHIA DeNot.

Length 128 μ; breadth 18 μ; apices 5.2 μ. Occasional, St. Anthony.

10. CLOSTERIUM ANGUSTATUM Kütz.

Length 310–560 μ; breadth 19–23 μ; apices 12 μ. Fairly common at St. Anthony.

Some of the specimens of this species show very interesting conditions of the surface ornamentation. One specimen, for instance, was divided into five nearly equal portions. The central portion was somewhat tumid, a character noted in many of the specimens from this material. The striae on this portion were very distinctly twisted laterally. The end segments show, even more distinctly, the twisting, subspiral arrangement of the striae. This twisting is developed more fully as the ends are approached. The intermediate areas of the membrane which were formed later by division do not keep to the subspiral striae, but have them in the normal, straight form. Nearly all the specimens from this locality show more or less of the subspiral arrangement of the costae.

11. CLOSTERIUM STRIOLATUM Ehrenb.

Length 275 μ; breadth 27 μ; apex 12 μ. Common, St. Anthony.

12. CLOSTERIUM PARVULUM Näg.

Length 115 μ; breadth 10 μ. Common, St. Anthony.

13. CLOSTERIUM TOXON West.

Length 290–325 μ; breadth 9–9.5 μ; apices 6 μ. Fairly common, St. Anthony.

DOCIDIUM Bréb. 1844

14. DOCIDIUM BACULUM Bréb.

Length 340 μ; breadth 21 μ; apex 13.3 μ. Common, St. Anthony.

PLEUROTAENIUM Näg. 1849

15. PLEUROTAENIUM CORONATUM (Bréb.) Rabenh.

Length 558 μ; breadth 60 μ; apex 42 μ. Common, Bay of Islands.

16. PLEUROTAENIUM EHRENBERGII (Bréb.) DeBary.

Length 405 μ; breadth 30 μ; apex 22 μ. Occasional, St. Anthony.

TETMEMORUS Ralfs. 1844

17. TETMEMORUS BREBISSONII TURGIDUS Ralfs.

Length 174–198 μ; breadth 40–44 μ. Abundant, Rose au Rue.

TETMEMORUS BREBISSONII MINOR DeBary.

Length 78 μ; breadth 18 μ; isthmus 13.5 μ. Common, St. Anthony.

TETMEMORUS BREBISSONII MINIMUS W. & G. S. West.

Length 62–65 μ; breadth 15–16 μ. Abundant, Rose au Rue.

On second examination, the form from this locality, reported in the previous paper as *T. minutus* DeBary, should be placed here.

18. TETMEMORUS GRANULATUS (Bréb.) Ralfs.

Length 190–204 μ; breadth 42–53 μ. Common at Bay of Islands and St. Anthony.

19. TETMEMORUS LAEVIS (Kütz.) Ralfs.

Length 133 μ; breadth 25 μ. Common, Rose au Rue.

EUASTRUM Ehrenb. 1832

20. EUASTRUM CRASSUM (Bréb.) Kütz.

Length 170 μ; breadth 85 μ; isthmus 21 μ; breadth of polar lobe 45 μ. Fairly common, St. Anthony.

21. EUASTRUM PINNATUM Ralfs.

Length 120–126 μ; breadth 57–60 μ; isthmus 16–18 μ; polar lobe 31–33 μ. Common, St. Anthony.

22. EUASTRUM OBLONGUM CEPHALOPHORUM West.

Length 170 μ; breadth 105 μ; isthmus 26 μ; polar lobe 75 μ. Occasional, Bay of Islands.

23. EUASTRUM AMPULLACEUM Ralfs.

Length 87–110 μ; breadth 47–72 μ; isthmus 15–20 μ; polar lobe 22–31 μ. Rose au Rue, Bay of Islands, and St. Anthony.

This is the most common species of the genus as far as the material examined shows. It was subject to little variation in form.

24. EUASTRUM OBESUM Josh.

Length 62 μ; breadth 34 μ; isthmus 8.2 μ; polar lobe 16 μ. Rare, St. Anthony. This is the first record for North America for this species. It was typical in its form and size.

25. EUASTRUM DIVARICATUM Lund.

Length 43 μ; breadth 37 μ; isthmus 6 μ; polar lobe 18 μ. Occasional, St. Anthony.

26. EUASTRUM ELEGANS (Bréb.) Kütz.

Length 25 μ; breadth 15 μ; isthmus 4 μ. Rare, Rose au Rue.

27. EUASTRUM BINALE (Turp.) Ehrenb.

Length 18 μ; breadth 15 μ. Rare, Rose au Rue.

28. EUASTRUM DENTICULATUM (Kirchn.) Gay.

Length 31 μ; breadth 23 μ; isthmus 5 μ; polar lobe 18 μ. Common, St. Anthony.

29. EUASTRUM PECTINATUM INEVOLUTUM W. & G. S. West.

Length 60–62 μ; breadth 42–44 μ; isthmus 9–12 μ; polar lobe 23–25 μ; thickness 25–27 μ. Very common, St. Anthony. This is the first North American record for this variety.

30. EUASTRUM FISSUM AMERICANUM Cushman.

Length 38 μ; breadth 23 μ; isthmus 6 μ; polar lobe 16 μ. Occasional, St. Anthony.

31. EUASTRUM ALLENII Cushman.

Length 105 μ; breadth 49 μ; isthmus 18 μ; polar lobe 33 μ. Occasional, Rose au Rue.

MICRASTERIAS Ag. 1827

32. MICRASTERIAS PINNATIFIDA (Kütz.) Ralfs.

Length 66 μ; breadth 75 μ; apical lobe 56 μ. Occasional, Bay of Islands.

33. MICRASTERIAS TRUNCATA (Corda) Bréb.

Length 78 μ; breadth 78 μ; isthmus 16 μ; apical lobe 56 μ. Common, St. Anthony.

34. MICRASTERIAS CONFERTA NOVAE-TERRAE Cushman.

Length 102 μ; breadth 93 μ; isthmus 12 μ; apical lobe 37 μ. Frequent, Rose au Rue.

35. MICRASTERIAS DENTICULATA Bréb.

Length 205 μ; breadth 170 μ; isthmus 30 μ: apical lobe 102 μ. Occasional, Bay of Islands.

MICRASTERIAS DENTICULATA ANGULOSA (Hantzsch.) W. & G. S. West.

Length 330 μ; breadth 210 μ; isthmus 36 μ; apical lobe 58 μ. Occasional, St. Anthony.

COSMARIUM Corda. 1834

36. COSMARIUM SUBUNDULATUM Wille.

Length 58 μ; breadth 41 μ; isthmus 16 μ. Common, St. Anthony.

37. COSMARIUM CUCUMIS (Corda) Ralfs.

Length 70–75 μ; breadth 39–42 μ; isthmus 19–21 μ. Common, St. Anthony.

38. COSMARIUM CONTRACTUM JACOBSENII (Roy) W. & G. S. West.

Length 43 μ; breadth 24 μ; isthmus 6 μ. Occasional, St. Anthony.

COSMARIUM CONTRACTUM ELLIPSOIDEUM (Elfv.) W. & G. S. West.
Length 33.5 μ; breadth 27 μ; isthmus 8 μ. Common, St. Anthony.

39. COSMARIUM HAMMERI Reinsch.
Length 41 μ; breadth 32 μ; isthmus 12 μ. Occasional, Bay of Islands.

40. COSMARIUM NYMANNIANUM Grun.
Length 43 μ; breadth 20 μ; isthmus 8 μ. Common, St. Anthony.

41. COSMARIUM GRANATUM Bréb.
Length 27 μ; breadth 20 μ; isthmus 7 μ. Common, St. Anthony.

42. COSMARIUM PSEUDONITIDULUM Nordst.
Length 53 μ; breadth 42 μ; isthmus 12 μ. Occasional, Bay of Islands.

43. COSMARIUM PYRAMIDATUM Bréb.
Length 87 μ; breadth 56 μ; isthmus 19 μ. Common, St. Anthony.

44. COSMARIUM PSEUDOPYRAMIDATUM Lund.
Length 50–56 μ; breadth 31–33 μ; isthmus 9–10 μ. Very common, Rose au Rue.

45. COSMARIUM VENUSTUM (Bréb.) Arch.
Length 37 μ; breadth 26 μ; isthmus 6.5 μ; apex 15 μ. Common, St. Anthony.

46. COSMARIUM CONNATUM Bréb.
Length 72 μ; breadth 56 μ; isthmus 45 μ. Rare, St. Anthony.

47. COSMARIUM IMPRESSULUM MINOR Turner.
Length 16 μ; breadth 12 μ; isthmus 3.5 μ. Common, Rose au Rue.

48. COSMARIUM SUBSPECIOSUM Nordst.
Length 50 μ; breadth 40 μ; isthmus 18.5 μ. Occasional, St. Anthony.

XANTHIDIUM Ehrenb. 1834

49. XANTHIDIUM ARMATUM (Bréb.) Rabenh.

Length without spines 148 μ; breadth without spines 93 μ, with spines 115 μ; isthmus 45 μ. Fairly common, Rose au Rue.

This species was also common in the material from St. Anthony, but was very much smaller than in that from Rose au Rue. The spines were for the most part simple, those with bifurcate spines being the exception. A considerable variation was noticed in the specimens from Rose au Rue. An average specimen from St. Anthony measures: length without spines 96 μ; breadth without spines 67 μ; with spines 80 μ; isthmus 32 μ.

50. XANTHIDIUM ANTILOPAEUM (Bréb.) Kütz.

Length with spines 96 μ; without 58 μ; breadth with spines 80 μ, without 46 μ; isthmus 15 μ. Occasional, St. Anthony.

XANTHIDIUM ANTILOPAEUM MINNEAPOLIENSE Wolle.

Length with spines 87 μ, without 68 μ; breadth with spines 75 μ; without 53 μ. Occasional, St. Anthony.

The specimens of this variety had the lateral spines below the chain of granules. The cells in the variety were larger than in the preceding but the lateral spines were very much shorter.

51. XANTHIDIUM CRISTATUM Bréb.

Length with spines 72 μ; breadth with spines 56 μ, without spines 40 μ; isthmus 12 μ. Common, St. Anthony.

ARTHRODESMUS Ehrenb. 1838

52. ARTHRODESMUS CONVERGENS Ehrenb.

Length with spines 45 μ, without 32 μ, breadth with spines 45 μ, without 28 μ; isthmus 6.5 μ. Common, St. Anthony.

53. ARTHRODESMUS QUADRIDENS Wood.

Length 34 μ; breadth with spines 50 μ, without 34 μ; isthmus 9 μ. Common, St. Anthony.

54. ARTHRODESMUS INCUS RALFSII W. & G. S. West.

Length 24 μ; breadth with spines 40 μ, without 19 μ; isthmus 7 μ. Common, St. Anthony.

ARTHRODESMUS INCUS LONGISPINUS Eichl. & Racib.

Length with spines 46 μ, without 26 μ; breadth with spines 34 μ, without 14 μ; length of spines 15 μ; isthmus 5 μ. Occasional, St. Anthony.

STAURASTRUM Meyen. 1829

55. STAURASTRUM ARISTIFERUM Ralfs.

Length 40 μ; breadth without spines 68 μ; isthmus 9 μ. Occasional, St. Anthony.

56. STAURASTRUM DUBIUM West.

Length 28 μ; breadth 22 μ; isthmus 10 μ. Very common, Rose au Rue.

57. STAURASTRUM LONGISPINUM BIDENTATUM (Wittr.) Cushman.

Length 90 μ; breadth with spines 108 μ, without 83 μ; isthmus 40 μ. Rare, St. Anthony.

58. STAURASTRUM ARCUATUM Nordst.

Breadth with processes 45–50 μ; isthmus 8.5 μ. Occasional, St. Anthony.

59. STAURASTRUM NANUM Wolle.

Length 25 μ; breadth with spines 31 μ, without 22 μ; isthmus 8 μ. Common, Rose au Rue.

60. STAURASTRUM JOHNSONII W. & G. S. West.

Length 56 μ; breadth 84 μ; isthmus 9.5 μ. Occasional, St. Anthony.

61. STAURASTRUM GRACILE Ralfs.

Length with processes 58 μ, without 15 μ; isthmus 8 μ. Not common, St. Anthony.

62. STAURASTRUM CRENULATUM (Näg.) Delp.

Breadth 25–37 μ; isthmus 9 μ. Rare, Rose au Rue. Frequent, St. Anthony.

63. STAURASTRUM MACROCERUM Wolle.

Length 48 μ; breadth with processes 100 μ; without 28 μ; isthmus 18 μ. Common, St. Anthony.

64. STAURASTRUM ECHINATUM Bréb.

Length without spines 45 μ; breadth with spines 50 μ, without 37 μ; isthmus 16 μ. Occasional, St. Anthony.

65. STAURASTRUM EUSTEPHANUM (Ehrenb.) Ralfs.

Breadth with processes 58 μ; isthmus 13 μ. Occasional, St. Anthony.

66. STAURASTRUM ARCTISCON (Ehrenb.) Lund.

Breadth with processes 100 μ, without 50 μ; isthmus 25 μ. Rare, St. Anthony.

SPHAEROZOSMA Corda. 1835

67. SPHAEROZOSMA EXCAVATUM Ralfs.

Breadth of filament 14 μ; length of cells 12 μ; isthmus 6.5μ. Rare, St. Anthony.

DESMIDIUM Ag. 1824

68. DESMIDIUM SWARTZII Ag.

Breadth of filament 34 μ; length of cell 18 μ. Rare, St. Anthony.

69. DESMIDIUM COARCTATUM Nordst.

Maximum breadth of filament 43.5 μ, minimum 31 μ; length 26 μ. Rare, St. Anthony.

70. DESMIDIUM QUADRATUM Nordst.

Breadth of filament 27–29 μ; length of cell 21–23 μ. Rare, St. Anthony.

71. DESMIDIUM BAILEYI (Ralfs) DeBary.

Breadth of filament 23–25 μ. Rare, Rose au Rue and St. Anthony.

GYMNOZYGA Ehrenb. 1840

72. GYMNOZYGA MONILIFORMIS Ehrenb.

Breadth of filament 22 μ; length of cell 28 μ. Occasional, Bay of Islands and St. Anthony.

BOSTON SOCIETY OF NATURAL HISTORY.

INDEX TO AMERICAN BOTANICAL LITERATURE
(1906)

The aim of this Index is to include all current botanical literature written by Americans, published in America, or based upon American material; the word America being used in its broadest sense.

Reviews, and papers which relate exclusively to forestry, agriculture, horticulture, manufactured products of vegetable origin, or laboratory methods are not included, and no attempt is made to index the literature of bacteriology. An occasional exception is made in favor of some paper appearing in an American periodical which is devoted wholly to botany. Reprints are not mentioned unless they differ from the original in some important particular. If users of the Index will call the attention of the editor to errors or omissions their kindness will be appreciated.

This Index is reprinted monthly on cards, and furnished in this form to subscribers, at the rate of one cent for each card. Selections of cards are not permitted; each subscriber must take all cards published during the term of his subscription. Correspondence relating to the card-issue should be addressed to the Treasurer of the Torrey Botanical Club

Arthur, J. C. Cultures of *Uredineae* in 1905. Jour. Myc. **12**: 11–27. 8 Mr 1906.

Atkinson, G. F. A new *Entoloma* from central Ohio. Jour. Myc. **12**: 236, 237. *pl. 92.* 5 D 1906.

E. subcostatum sp. nov.

Berry, E. W. Leaf-rafts and fossil leaves. Torreya **6**: 246–248. *f. 2.* 13 D 1906.

Bessey, C. E. Botanical notes. Science II. **24**: 571, 572. 2 N 1906.

Includes a teratological note on *Cassia Chamaecrista.*

Blakeslee, A. F. Zygospores and sexual strains in the common bread mould, *Rhizopus nigricans.* Science II. **24**: 118–122. 27 Jl 1906.

Blanchard, W. H. A new dwarf blackberry. Torreya **6**: 235–237. 26 N 1906.

Rubus abbrevians sp. nov., from Vermont.

Borge, O. Süsswasser-Chlorophyceen von Feuerland und Isla Desolacion. Bot. Stud. Kjellman 21–34. *pl. 2.* 1906.

Includes descriptions of new species in *Oocystis, Cosmarium* (2), *Penium,* and *Closterium.*

Burnham, S. H. A new species of *Monotropsis.* Torreya **6**: 234, 235. 26 N 1906.

M. Lehmanae.

Conzatti, C. Taxinomia de las Orquídeas mexicanas (fin). Mem. Soc. Ci. Ant. Alzate **21**: 273–341. "1904." [1906.]

Cook, M. T. Gall-insects and insect-galls. Science II. **24**: 312. 7 S 1906.

Cook, M. T. Herbarium type specimens in plant morphology. Science II. **24**: 217, 218. 17 Au 1906.

Dowell, P. North American species of *Calceolaria*. Bull. Torrey Club **33**: 547–556. *pl. 18-22*. 19 D 1906.
Enumerates 16 species, of which 7 are new.

Durand, E. J. *Peziza fusicarpa* Ger. and *Peziza semitosta* B. & C. Jour. Myc. **12**: 28–32. 8 Mr 1906.

Fairchild, D. G. Cats as plant investigators. Science II. **24**: 498, 499. 19 O 1906.

Foslie, M. & Howe, M. A. Two new coralline algae from Culebra, Porto Rico. Bull. Torrey Club **33**: 577–580. *pl. 23–26*. 19 D 1906.
Goniolithon acropetum sp. nov., and *Lithophyllum Antillarum* sp. nov.

Fries, R. E. Morphologisch-anatomische Notizen über zwei südamerikanischen Lianen. Bot. Stud. Kjellman 89–101. *f. 1–4*. 1906.
Cissus sp. and *Siolmatra brasiliensis*.

Gager, C. S. De Vries and his critics. Science II. **24**: 81–89. 20 Jl 1906.

Giacomelli, E. Observaciones é investigaciones sobre la *Prosopanche Burmeisteri* De Bary. Anal. Soc. Ci. Argent. **62**: 5–22. Jl 1906.

Harper, R. M. Some hitherto undescribed outcrops of Altamaha Grit and their vegetation. Torreya **6**: 241–246. *f. 1*. 13 D 1906.

Hill, E. J. A Mississippi *Aletris* and some associated plants. Torreya **6**: 231, 232. 26 N 1906.

Howard, B. J. Tannin cells of persimmons. Bull. Torrey Club **33**: 567–576. *f. 1–8*. 19 D 1906.

Howe, M. A. Some photographs of the silk-cotton tree (*Ceiba pentandra*), with remarks on the early records of its occurrence in America. Torreya **6**: 217–231. *f. 1–6*. 26 N 1906.

Howe, R. H. Lichens of Mount Monadnock, New Hampshire. Am. Nat. **40**: 661–665. 21 S 1906.

Jordan, D. S. Discontinuous variation and pedigree culture. Science II. **24**: 399, 400. 28 S 1906.

Kellerman, W. A. Fungi selecti guatemalenses. Exsiccati decade I. Jour. Myc. **12**: 238–241. 5 D 1906.

Kellerman, W. A. Uredineous culture experiments with *Puccinia Sorghi*, 1905. Jour. Myc. **12**: 9–11. 8 Mr 1906.

Kellogg, V. L. Scientific aspects of Luther Burbank's work. Pop. Sci. Mo. **69**: 363–374. O 1906. [Illust.]

Lindman, C. A. M. Zur Kenntnis der Corona einiger Passifloren. Bot. Stud. Kjellman 55–79. *f. 1–12.* 1906.
Based upon studies in Brazil and Paraguay, in 1892–1894.

Long, W. H. Notes on new or rare species of Ravenelia. Jour. Myc. **12**: 233–236. 5 D 1906.
Includes descriptions of 2 new species.

MacDougal D. T. Discontinuous variation. Science II. **24**: 730, 731. 7 D 1906.

MacDougal, D. T. Discontinuous variation in pedigree-cultures. Pop. Sci. Mo. **69**: 207–225. *f. 1–8.* S 1906.

Maza, M. G. de la. Determinacion de plantas cubanas (fanerógamas). Revista Facult. Let. & Ci. Univ. Habana **2**: 55–67. Ja 1906.

Maza, M. G. de la. Formas interesantes de reino vegetal. Revista Facult. Let. & Ci. Univ. Habana **2**: 128–139. Mr 1906.

Morgan, A. P. North American species of *Lepiota*. Jour. Myc. **12**: 154–159. 29 S 1906; 195–203. 8 N 1906; 242–258. 5 D 1906.
Includes descriptions of 9 new species.

Morgan, A. P. North American species of *Marasmius* (continued). Jour. Myc. **12**: 1–9. 8 Mr 1906.
Includes a description of *M. felix* sp. nov.

Murrill, W. A. How Bresadola became a mycologist. Torreya **6**: 233, 234. 26 N 1906.

Oliver, F. W. Botany in England. Science II. **24**: 321–327. 14 S 1906.

Pammel, L. H. Northern limit of the papaw tree. Science II. **24**: 48. 13 Jl 1906.

Pulle, A. Ueber einige neue und seltene Arten aus Surinam. Recueil Trav. Bot. Néerl. **2**: 193–208. 1906.
Mere reprints of the diagnoses of the new species published a few months earlier in his Enumeration of the vascular plants known from Surinam.

Shafer, J. A. *Hibiscus oculiroseus* Britton. Torreya **6**: 233. 26 N 1906.

Sheldon, J. L. A rare *Uromyces*. Torreya **6**: 249, 250. 13 D 1906.

Skottsberg, C. Observations on the vegetation of the Antarctic sea. Bot. Stud. Kjellman 245–264. *pl. 7–10.* 1906.

White, D. [The flora of the Brazilian coal measures.] Science II. **24**: 378, 379. 21 S 1906.

Wildeman, E. de. *Atriplex confertifolia* Ser. Wats. Ic. Sel. Hort. Thenensis **5**: 139–141. *pl. 194.* F 1906.
Native of western North America.

Wildeman, E. de. *Dicliptera Niederleiniana* Lindau. Ic. Sel. Hort. Thenensis **5**: 153–155. *pl. 198.* Mr 1906.
Native of Argentina.

Wildeman, E. de. *Rhipsalis trigona* Pfeiff. Ic. Sel. Hort. Thenensis **5**: 135–137. *pl. 193.* F 1906.
Native of Brazil.

Wilson, G. W. The identity of *Mucor Mucedo.* Bull. Torrey Club **33**: 557–560. 19 D 1906.

Wooton, E. O. Southwestern localities visited by Charles Wright. Bull. Torrey Club **33**: 561–566. 19 D 1906.

INDEX TO VOLUME 33

New names, and the final members of new combinations, are in **bold face type.**

Abies, 405, 411, 514, 520; alba, 432; balsamea, 406, 412, 434, 436; canadensis, 434; concolor, 406; Douglasii 431; Fraseri, 406; grandis, 412, 430, 436; lasiocarpa, 412, 433, 435, 514; magnifica, 406; nigra, 426–428, 432; nobilis, 406; pectinata, 412, 432; religiosa, 412, 435; subalpina, 435

ABRAMS, L. R. Two new southwestern species of Pentstemon, 445

Absorption of atmospheric moisture by desert shrubs, 367

Abutilon incanum, 374

Acer, 69, 282, 284, 286, 288, 294; platanoides, 281, 304; Pseudo-Platanus, 281

Achillea **alpicola,** 157; lanulosa alpicola, 157

Aconitum Napellus, 334, 340

Acrostichum, 36, 197, 457, 464, 604; aculeatum, 41; aureum, 463, 464; axillare, 603, 604; buxifolium, 45; decrescens, 46; erythrodes, 595; excelsum, 463, 464, 470; excelsum f. **lobatum,** 464; gracile, 41; guineensis, 46; japurense, 598; laurifolium, 37; lomarioides, 47, 464; Meyerianum, 39; Pittieri, 600; scandens, 37, 39; sorbifolium, 35, 36, 44, 591, 592, 600, 602; spectabile, 47; tennifolium, 39

Actinella Brandegei, 156; glabra, 155; grandiflora glabrata, 156

Actinostachys, 462; Germani, 191

Adelia acuminata, 534

Adiantum, 272; clavatum, 200; Farleyense, 471; hispidulum, 192; modestum, 192; tenerum, 471

Aecidium, 404, 518, 521; abietinum, 430; **Argithamniae,** 33; balsameum, 435; **Cardui,** 33; carneum, 414; Cerebrum, 423; coloradense, 426; columnare, 432; conorum-Piceae, 431; decolorans, 428; deformans, 423; elatinum, 434; Engelmanni, 431; Ephedrae, 437; **Falcatae,** 32; filamentosum, 418; giganteum 423; **Herrerianum,** 520; Laricis, 436; ornamentale, 433; Peckii, 433; pseudo-balsameum, 430; **pseudo-columnare,** 436; **pulverulentum,** 521; pyriforme 419; Ravenelii, 414; **Sorbi,** 521; **Triostei,** 32

Agardhiella, 83

Agastache **pallidiflora,** 150

Agropyron spicatum, 29

Agrostemma, 490

Alabama plants, Notes on the distribution of some, 523

Alchemilla, 328, 492; arvensis, 340, 493

Algae from Culebra, Two new coralline, 577

Allium, 273, 278, 283; fistulosum, 298, 300

Alopecurus pratensis, 490

Alsinastrum, 138

Alsine, 435

Alsinopsis **obtusiloba,** 140; **propinqua,** 140; **quadrivalvis,** 140; **Rossii,** 140

Amarella **monantha,** 148; **plebeia Holmii,** 148

Amelanchier, 237, 310; alnifolia, 310; rotundifolia, 310; **Scudderi,** 310; typica, 310

American botanical literature, Index to (1901–1903), 207; (1905), 59, 263, 447; (1906), 125, 319, 353, 397, 505, 541, 581, 617

American ferns, 189, 591

American species of Stenochlaena, The, 591

American tropics, The species of Lycopodium of the, 101

Amphicarpa monoica, 32

Amsonia angustifolia, 241; ciliata, 240, 241; ciliata filifolia, 241; tenuifolia, 241

Amygdalus gracilis, 308

Anchistea, 528; virginica, 523, 527, 528

Androcera lobata, 150; **rostrata,** 150

Andromeda Novae-Caesareae, 181; Parlatorii, 181

Androsace septentrionalis subulifera, 148; **subulifera,** 148

Aneimia adiantifolia, 457, 461, 472

Aneura palmata, 129

Anisolotus, 144; **brachycarpus,** 144; **rigidus,** 144; **Wrightii,** 144

Annularia Fenzlii, 216; **sphaerospora,** 216

Anogra latifolia, 146, v

Anomalous anther-structure of Dicorynia, Duparquetia, and Strumpfia, 223

Anonymos, 242; caroliniensis, 241, 242

Anoplanthus, 151

Antennaria solitaria, 536

Anther-structure of Dicorynia, Duparquetia, and Strumpfia, 223
Anthoceros, 274, 299
Anthopogon **barbellatus,** 148; **elegans,** 148
Aphyllon, 151; ludovicianum, 151; multiflorum, 151
Apios Apios, 32; tuberosa, 32
Aplopappus, 152; Parryi, 153
Apocynum, 73; cannabinum, 73
Arabis georgiana, 532; longirostris, 142
Aragallus **Hallii,** 144
Aralia Daphnophyllum, 179
Arenaria biflora, 140; hirta, 140; obtusa, 140; propinqua, 140; quadrivalvis, 140; Rossii, 140; sajanensis, 140; verna aequicaulis, 140; verna hirta, 140
Argentina anserina concolor, 143; **argentea,** 143
Argithamnia Schiedeana, 34
Arisaema, 273
Artemisia cana viscidula, 157; desertorum Scouleriana, 157; **pabularis,** 157; rhizomata pabularis, 157; **Scouleriana,** 157; **viscidula,** 157
Arthrobotrya, 40
Arthrodesmus, 613; convergens, 613; Incus longispinus, 614; Incus Ralfsii, 613; quadridens, 613
ARTHUR, J. C. New species of Uredineae, 27, 513
ARTHUR, J. C. & KERN, F. D. North American species of Peridermium, 403
Arum, 328
Asclepias, 74; incarnata, 71, 72, 76; syriaca, 73
Ascophora Mucedo, 559, 560
Aspidium coriandrifolium, 200; fimbriatum, 199; heracleifolium, 200; stipularis, 198; trifoliatum, 481
Asplenium, 189, 193; Adiantum-nigrum, 193; anchorita, 194; Andrewsii, 193; auritum, 196; biscaynianum, 195, 473, 474; **Curtissii,** 194; **dentatum,** 195, 473, 474; erosum, 196; montanum, 193; monteverdense, 476; muticum, 195; myriophyllum, 195, 476; platyneuron Hortonae, 190; rhizophyllum, 195; rhizophyllum biscaynianum, 195, 473; serratum, 475; serratum f. **incisum,** 475; **verecundum,** 193, 195, 473–475
Aster **polycephalus,** 153; scoparius, 153
Atlantic coastal plain, Mesozoic flora of the, 163
Atmospheric moisture: its absorption by desert shrubs, 367
Atriplex, 374; patula subspicata, 137; **subspicata,** 137
Avena, 219
Azalea, 239; arborescens, 239, 240; lutea, 239
Azolla caroliniana, 484

Bahamas, Vernonia in the, 183
Bahia, 154, 156; nudicaulis, 155; oblongifolia, 155
Baldwinia atropurpurea, 237
Baptisia bracteata, 533
Bartonia, 240; lanceolata, 240; tenella brachiata, 240
Batatas, 495; pentaphyllus, 502
Batschia, 242; carolinensis, 241, 242
Behavior of the pollen-tube in Houstonia coerulea, The, 487
Berberis Aquifolium, 141; Fremontii, 141; nana, 141; nervosa, 141; repens, 141
Bermuda, Hepaticae of, 129
BERRY, E. W. Contributions to the Mesozoic flora of the Atlantic coastal plain, 163
Berthelotia, 154; **sericea,** 154
Betula, 329, 437; alba, 493
Bigelovia Douglasii latifolia, 152; Wrightii, 152
Bistorta, 57; **bistortoides,** 57
Blechnum serrulatum, 472
Bolbitis, 604
Bolophyta alpina, 156
Botanical literature, American, (1901–1903), 207; (1905), 59, 263, 447; (1906), 125, 319, 353, 397, 505, 541, 581, 617
Botrychium, 189, 274, 280, 292, 293, 298; alabamense, 190; californicum, 190; obliquum, 191, 292, 306; occidentale, 190; onondagense, 191; salaifolium, 190; tennifolium, 191; ternatum, 292
Brachyphyllum macrocarpum, 168, 182
Brittonastrum Greenei, 150; pallidiflorum, 150
Brunnichia cirrhosa, 534
Bryonia, 330, 332
Bubakia, 520

Cactus similis, 146, v
Caeoma, 404, 520; Abietis-canadensis, 404; conigenum, 405, 520; Laricis, 404; **strobilina,** 519
Cafraria, 37, 39
Calamintha, 244, 245; caroliniana, 243, 245; grandiflora, 243, 244
Calceolaria, 547, 548; **angustifolia,** 549, 554, 556; **brevis,** 548, 552, 556; fruticulosa, 548, 550; fruticulosa **flavescens,** 551; **glabra,** 548, 552, 556; glutinosa, 548, 549; **humilis,** 549, 555; **longipes,** 548, 551, 556; **nigricans,** 549, 554; procumbens,

548, 551; riparia, 549, 554; riparia **Houstoni,** 554; **Rosei,** 549, 555, 556; stricta, 549, 553, 556; **tenuifolia,** 548, 550, 556; verbenacea, 549, 553; verticillata, 548, 549, 556
Calceolaria, North American species of, 547
Callithamnion, 83
Calochortus, 537–540; apiculatus, 539; ciliatus, 539; elegans, 537, 538, 540; elegans Lobbii, 539; elegans major, 540; elegans minor, 540; elegans nanus, 537; Lobbii, 539; Lyallii, 537, 539, 540; **nanus,** 537; Purdyi, 538; subalpinus, 538
Calochortus, Notes on, 537
Calocylindrus diplospora, 348
Calonyction muricatum, 318; speciosum muricatum, 318; **tastense,** 318
Calyptospora, 405–407, 432; Goeppertiana, 432
Campanula, 405; americana, 417
Campyloneurum, 192; angustifolium, 192, 467; costatum, 192, 467; latum, 192, 468; Phyllitidis, 192, 458, 467–469
Canavalia ensiformis, 31
CARDIFF, I. D. A study of synapsis and reduction, 271
Carduus arvensis, 33; **filipendulus,** 157; **florissantensis,** 311, 312; Hookerianus, 33; lanceolatus, 33
Carex, 439–443; **agglomerata,** 442, 443; Backii, 439; cephaloidea, 442, 443; cephalophora, 441; Clarkii, 169; concinna, 440; **concinnoides,** 440; durifolia, 439; gravida, 441, 442; Leavenworthii, 442; **mediterranea,** 441, 442; Muhlenbergii, 441–443; picta, 526, 527, 536; Richardsonii, 440; **saximontana,** 439
Carex, Notes on, 439
Carpinus, 328
Carpites Liriophylli, 174
Carya, 328; olivaeformis, 339
Celtis pallida, 372
Cephalozia connivens, 129–131; divaricata, 130, 131
Cerastium, 435
Cerasus demissa, 143; demissa melanocarpa, 143
Ceratelium (see **Cerotelium**), 30; **Canavaliae,** 30
Ceratolejeunea, 3
Ceratonia Siliqua, 567
Cereus conoideus, 146; phoeniceus, 146; Roemeri, 146
Ceropteris viscosa, 196
Cerotelium, 514
Chaenactis scaposa, 156
Chaetochloa macrosperma, 519
Chamaechaenactis, 155; **scaposa,** 156
Chamaesyce **albicaulis,** 145; **flagelliformis,** 144; **rugulosa,** 145
Changes accompanying secretion in the nectar-glands of Vicia Faba, 247
Cheilanthes, 193
Cheilolejeunea, 1, 2, 4, 5, 8, 9, 16, 17, 19; **aneogyna,** 2, 3, 7,; decidua, 4, 6, 7, 25; emarginuliflora, 14; heteroclada, 2; jamaicensis, 7, 17; lineata, 3, 4, 17; phyllobola, 3; versifolia, 3, 12, 13
Cheiranthus argillosus, 141; aridus, 141; asperrimus, 141; Bakeri, 141; nivalis amoenus 142
Chenopodium leptophyllum oblongifolium, 137; **oblongifolium,** 137; subspicatum, 137
Chrysomyxa, 405–407; Abietis, 518; Ledi, 429, 430; Pyrolae, 432
Chrysopsis graminifolia, 536; oligantha, 536
Chrysothamnus glaucus, 152; **latifolius,** 152; **serrulatus,** 152
Cibotium Barometz, 478
Cinnamomum Heerii, 179; intermedium, 179, 182
Circaea lutetiana, 533
Cirsium filipendulum, 157; Hookerianum, 33; virginianum filipendulum, 157
Cissites crispus, 177
Citrullus, 332
Claytonia, 138–140; arenicola, 138; asarifolia, 138; Chamissoi, 139; parviflora, 139; parviflora depressa, 139; parvifolia, 139; perfoliata, 139; sibirica, 138; triphylla, 140; virginica, 281
Cleome Sonorae, 142
Clinopodium, 244, 245; carolinianum, 243–245; **georgianum,** 243; grandiflorum, 244; Walterianum, 244
Clitoria cajanifolia, 515
Closterium, 608; angustatum, 608; Cynthia, 608; Digitus, 348; parvulum, 609; striolatum, 609; toxon, 609
Cnicus altissimus filipendulus, 157
Coastal plain, Mesozoic flora of the Atlantic, 163
Coastal plain of Georgia, Some new or otherwise noteworthy plants from the, 229
COCKERELL, T. D. A. Fossil plants from Florissant, Colorado, 307
Coleosporium, 405, 407, 408, 417; Campanulae, 405, 417; **Eupatorii,** 31; Senecionis minus, 437; Steviae, 31
Collybia **brunnescens,** 214

Cololejeunea Jooriana, 130, 131; minutissima, 131
Color in plants, 77
Colorado, Fossil plants from, 307
Colutea, 175; antiqua, 176; Boweniana, 176; coronilloides, 175; debilis, 176; Langeana, 175; macrophylla, 176; **obovata,** 175, 182; primordialis, 175, 176; protogaea, 175; Salteri, 176; valde-inaequalis, 176
Comptonia asplenifolia, 29; peregrina, 29
Contributions to the Mesozoic flora of the Atlantic coastal plain, 163
Convallaria, 278, 283; majalis, 298
Convolvulaceae, Studies in the North American, 313, 495
Convolvulus alatus, 497; bicolor, 318; copticus, 500; dissectus, 500; gossypiifolius, 500; macrocarpus, 496, 499; muricatus, 318; nemorosus, 502; operculatus, 499; pentaphyllus, 502; sublobatus, 318; tamnifolius, 314; triquetra, 497; tuberosus, 500; ventricosus, 498
Conyza, 153; cinerea, 184; Coulteri, 154; squarrosa, 153
Conyzella, 153, 154
Coralline algae from Culebra, Two new, 577
Cornella, 147; **canadensis,** 147; **suecica,** 147; **unalaschkensis,** 147
Cornus, 69, 147; canadensis, 147; florida, 147; mas, 147; mascula, 147; Nuttallii, 147; sanguinea, 147; sessilis, 147; suecica, 147; unalaschkensis, 147
Corylus, 328
Cosmarium, 611; connatum, 612; contractum ellipsoideum, 612; contractum Jacobsenii, 611; Cucumis, 611; granatum, 612; Hammeri, 612; impressulum minor, 612; Nymannianum, 612; pseudonitidulum, 612; pseudopyramidatum, 612; pyramidatum, 612; subspeciosum, 612; subundulatum, 611; venustum, 612
Covillea, 373, 375; tridentata, 373
Crataegus acerifolia, 311; **Lesquereuxii,** 311
Craterellus **Pogonati,** 218
Cressa, 315; aphylla, 315; cretica, 315; depressa, 315; **insularis,** 315; truxillensis, 315
Crocus, 328
Cronartium, 31, 405, 407, 424; **Comptoniae,** 29; ribicola, 425
Crossotolejeunea, 134; **bermudiana,** 132, 134–135; Boryana, 134; paucispina, 134
Croton, 520
Crotonophyllum cretaceum, 172
Crunocallis, 139; **Chamissonis,** 139
Cryptogramme Stelleri, 30
Cucumis, 330
Cucurbita, 328, 330–333, 337, 340; maxima, 329; Pepo, 330–333, 336, 338
Cucurbitaceae, The pollen-tube in some of the, 327
Culebra, Two new coralline algae from, 577
Cunninghamites squamosus, 165
CUSHMAN, J. A. New England desmids of the sub-family Saccodermæ, 343; some desmids from Newfoundland, 607
Cuttings of ivy, Influence of moisture upon the formation of roots by, 93
Cyclanthera, 327, 331–336, 338, 342; explodens, 331, 338
Cyclolejeunea, 13
Cylindrocystis, 347, 350, 351, 607; americana, 351; americana minor, 348, 607; Brebissonii, 347, 351, 607; Brebissonii minor, 347; crassa, 347, 351; diplospora, 348, 351, 608; diplospora **minor,** 348
Cynomorium, 492
Cynoxylon, 147
Cyrtolejeunea, 3
Cystolejeunea, 1, 4, 16, 17; **lineata,** 17, 19, 25
Cystopteris, 30
Cytological changes accompanying secretion in the nectar-glands of Vicia Faba, 247

Daphnogene Heerii, 179
Dasya, 83
Dasystephana **affinis,** 149; **Bigelovii,** 149; **Forwoodii,** 149; **interrupta,** 149; **Parryi,** 149; **Romanzovii,** 148
Davallia, 483; achillaefolia, 40; clavata, 200
Delphinium occidentale reticulatum, 140; **reticulatum,** 140
Dennstaedtia, 193
Desert shrubs, Absorption of atmospheric moisture by, 367
Desmidium, 615; Baileyi, 615; coarctatum, 615; quadratum, 615; Swartzii, 615
Desmids from Newfoundland, 607
Desmids of the sub-family Saccodermae, New England, 343
Dichromena ciliata, 32; radicans, 32
Dicorynia, 223, 224
Didymoglossum, 201; Krausii, 201; sphenoides, 201
Dietelia, 31
Digitalis, 337

Diodia, 487, 489; teres, 487; virginiana, 487
Diospyros rotundifolia, 181
Diphasium, 102, 120
Diplopappus ericaefolius hirtellus, 153
Distegia involucrata, 152, v; nutans, 152
Distribution of some Alabama plants, Notes on the, 523
Docidium, 609; Baculum, 609
Dodecatheon radicatum sinuatum, 148; **sinuatum,** 148
Dolicholus texanus, 27
Dolichos, 513; reticulatus, 28
DOWELL, P. North American species of Calceolaria, 547
Dramia, 274
Drosera, 247, 258, 278, 279, 302; longifolia, 279; rotundifolia, 261, 279, 298
Drymocallis glandulosa, 219
Drynaria, 603
Dryopteris, 198, 469, 477; ampla, 198, 476; aquilonaris, 197; Boottii, 198; cristata Clintoniana, 476; floridana, 476, 527; fragrans, 197; marginalis, 197; noveboracense, 476; parasitica, 477; patens, 477–479; patens **glandulosa,** 477; pittsfordensis, 197; reptans, 477; reticulata, 198; setigera, 477; simulata, 477; spinulosa, 197, 198; **stipularis,** 198, 478; Thelypteris, 479; unita glabra, 479
Dumortiera hirsuta, 129, 130
Duparquetia, 223, 224, 226

EATON, A. A. Pteridophytes observed during three excursions into southern Florida, 455
Ecballium, 330, 332; Elaterium, 330
Echinocereus **aggregatus,** 146; coccineus, 146; **Roemeri,** 146
Eleocharis, 515; palustris, 514
Elodea, 334; canadensis, 341
Elytraria, 242
Encelia, 374; farinosa, 374
Ephedra, 405, 412; antisyphilitica, 437; californica, 412, 438; nevadensis, 412, 437; oxycarpa, 437; pedunculata, 412, 437; Torreyana, 412, 438; trifurca, 412, 438
Epilobium **adenocladon,** 146; paniculatum adenocladon, 146
Equisetum, 190, 272, 301; Ferrissii, 201; Funstoni, 201
Erigeron, 153; cinereus, 153; **Wootonii,** 153
Eriocaulon lineare, 237, 527
Eriogonum aureum, 51; **coloradense,** 53; corymbosum, 55; corymbosum divaricatum, 55; divaricatum, 55; **divergens,** 55; effusum nudicaule, 54; **Fendlerianum,** 55; flavum, 51; **fusiforme,** 56; inflatum, 56; lachnogynum, 52, 53; lonchophyllum, 54, 55; microthecum Fendlerianum, 55; **nudicaule,** 54; ochroleucum, 53; **orthocaulon,** 53; ovalifolium, 53; pauciflorum, 54; salicinum, 55; **scoparium,** 54; **Tetraneuris,** 52; **tristichum,** 55; umbellatum, 52; **umbelliferum,** 51; villiflorum, 52; **xanthum,** 51
Eritrichium glomeratum hispidissimum, 150
Erocallis, 139; **triphylla,** 140
Erysimum **amoenum,** 142; **argillosum,** 141; **asperrimum,** 141; **Bakeri,** 141; nivale, 141; pumilum, 141; radicatum, 141
Erythraea **arizonica,** 148; calycosa arizonica, 148
Eschenbachia, 153, 154; **Coulteri,** 153
Etiolation, Studies in, 67
Euastrum, 610; Allenii, 611; ampullaceum, 610; binale, 610; crassum, 610; denticulatum, 610; divaricatum, 610; elegans, 610; fissum americanum, 611; obesum, 610; oblongum cephalophorum, 610; pectinatum inevolutum, 610; pinnatum, 610
Eucalyptus attenuata, 180; Geinitzi, 180; Wardiana, 180
Euclaytonia, 138
Euklisia, 142; **cordata,** 142; **crassifolia,** 142; hyacinthoides, 142; **longirostris,** 142
Eukrania, 147
Euosmolejeunea, 2–5, 16, 17; opaca, 132; trifaria, 5
Eupatorium macrophyllum, 31
Euphorbia, 69; albicaulis, 145; arkansana coloradensis, 145; corollata, 69, 70, 76; Cyparissias, 73; flagelliformis, 144; petaloidea flagelliformis, 144; serpyllifolia rugulosa, 145
Euschizaea, 462
Eustenochlaena, 37, 39
EVANS, A. W. Hepaticae of Puerto Rico, 1; The Hepaticae of Bermuda, 129
Evolvulus, 315; **adscendens,** 317; albiflorus, 317; discolor, 316; incanus, 317; microphyllus, 317; **Palmeri,** 317; **Wilcoxiana,** 315; **Wrightii,** 316
Excursions into southern Florida, Pteridophytes observed during three, 455

Falcata comosa, 32
Fallugia acuminata, 143, v; paradoxa acuminata, 143
Ferns, American, 189, 591

Ferns: species added to the flora of the United States from 1900 to 1905, 189
Ficus aligera, 172; crassipes, 172; daphnogenioides, 173, 182; Krausiana, 172; Woolsoni, 172
Fimbristylis Holwayana, 28; polymorpha, 28
Flabellaria magothiensis, 170
Flammula carbonaria, 217; **condensa,** 217; squalida, 217
Florida, Pteridophytes observed during three excursions into southern, 455
Florissant, Fossil plants from, 307
Formation of roots by cuttings of ivy. Influence of moisture upon the, 93
FOSLIE, M. & HOWE, M. A. Two new coralline algae from Culebra, Porto Rico, 577
Fossil plants from Florissant, Colorado, 307
Fouquieria, 370–372, 375; splendens, 367–371
Fragaria firma, 143; **ovalis,** 143
Franseria, 374; deltoidea, 374
Frasera **angustifolia,** 149; **scabra,** 149; speciosa angustifolia, 149; speciosa scabra, 149; speciosa stenosepala, 149; **stenosepala,** 149
Fraxinus caroliniana, 459
Frenelopsis, 168; gracilis, 167; Hoheneggeri, 168; leptoclada, 168
Frullania squarrosa, 132
Fungi, New species of, 213

Galium **subbiflorum,** 152; trifidum subbiflorum, 152
Galtonia candicans, 276
Gaylussacia frondosa, 238
Gentiana affinis, 149; barbellata, 148; Bigelovii, 149; elegans, 148; Forwoodii, 149; frigida, 149; interrupta, 149; monantha, 148; Moseleyi, 148; Parryi, 149; plebeia Holmii, 148; Romanzovii, 148; tenella, 148
Genus Operculina, The, 495
Genus Stenochlaena, The, 35, 591
Genus Vernonia in the Bahamas, The, 183
Georgia, Some new or otherwise noteworthy plants from the coastal plain of, 229
Gilia Nuttallii, 149; Watsoni, 149
Ginkgo, 289, 291; biloba, 289, 305
GLEASON, H. A. The genus Vernonia in the Bahamas, 183; The pedunculate species of Trillium, 387
Gleichenia Saundersii, 163; Zippei, 164
Gonatozygon, 344, 350; aculeatum, 345, 350; asperum, 344; Brebissonii, 344, 350; monotaenium, 344, 350; monotaenium **minutum,** 344; pilosum, 344, 350; Ralfsii, 344
Goniolithon **acropetum,** 577, 578, 580; frutescens, 578; frutescens f. flabelliformis, 578; frutescens f. typica, 578; strictum, 578
Goniopteris, 469
Grewiopsis flabellata, 177
Griffithsia, 83
Grinnellia, 83
Gymnoconia, 518
Gymnogramme triangularis viscosa, 196
Gymnopteris, 604; contaminoides, 605
Gymnozyga, 615; moniliformis, 615

Hamamelis, 329; virginiana, 341
Haplosporella, 219; **commixta,** 219
HARPER, R. M. Some new or otherwise noteworthy plants from the coastal plain of Georgia, 229; Notes on the distribution of some Alabama plants, 523
HARRIS, J. A. The anomalous anther-structure of Dicorynia, Duparquetia, and Strumpfia, 223
Hedera cretacea, 180; Helix, 93, 98
Helleborus, 278; foetidus, 298
Hepatica, 531; acutiloba, 531; triloba, 531
Hepaticae of Bermuda, 129
Hepaticae of Puerto Rico, 1
Hewittia, 318; bicolor, 318; sublobata, 318
Hicoria, 328
Hippeastrum, 328
Hosackia, 144; brachycarpa, 144; rigida, 144; Wrightii, 144
HOUSE, H. D. Studies in the North American Convolvulaceae, 313, 495
Houstonia, 488–493; coerulea, 487, 488
Houstonia coerulea, The behavior of the pollen-tube in, 487
HOWARD, B. J. Tannin cells of persimmons, 567
HOWE, M. A. & FOSLIE, M. Two new coralline algae from Culebra, Porto Rico, 577
Hyalopsora **pellaeicola,** 30
Hybanthus, 547
Hydnum **Blackfordae,** 218
Hydrophora stercorea, 560
Hygrolejeunea, 17
Hygrophorus **Davisii,** 214; **mephiticus,** 213
Hymenaea dakotana, 176
Hymenoxys, 156; chrysanthemoides multiflora, 157; chrysanthemoides Osterhoutii, 157; **macrantha,** 156; **multiflora,** 157; multiflora **Osterhoutii,** 157; **pumila,** 156; Richardsonii macrantha, 156; Richardsonii pumila, 156
Hypolepis repens, 192

Identity of Mucor Mucedo, The, 557
Ilex, 309; **Leonis,** 311; quercifolia, 311; strangulata, 177
Illicium, 535; floridanum, 530, 535
Index to American botanical literature (1901–1903), 207; (1905), 59, 263, 447; (1906), 125, 319, 353, 397, 505, 541, 581, 617
Influence of moisture upon the formation of roots by cuttings of ivy, 93
Inocybe **desquamans,** 216; **Sterlingii,** 217; vatricosa, 217
Inula, 153
Ionidium, 547, 548; angustifolium, 554; fruticulosum, 550; fruticulosum dentatum, 550, 551; glutinosum, 549; lineare, 549, 556; parietariaefolium, 554; parietariaefolium Houstoni, 554; polygalaefolium, 547, 549; procumbens, 551; riparium, 554; stipulaceum, 549; strictum, 553; verbenaceum, 553
Ipomoea, 495, 496; aegyptia, 502; alata, 498; alatipes, 499; altissima, 497; alulata, 497; ampliata, 503; bona-nox purpurascens, 318; codonantha, 503; coptica, 500; dissecta, 500; eriocephala, 314; Hamiltoni, 497; kentrocaulos, 500; lacioclados, 313; montana, 314; muricata, 318; nigricans, 500; nuda, 500; operculata, 499; Palmeri, 502; Palmeri platyphylla, 502; pentaphylla, 502; pilosa, 502; pterodes, 497, 499; rhodocalyx, 498; serrata, 314; sinuata, 500; spinulosa, 318; subpedata, 500; tamnifolia, 314; tastensis, 318; triquetra, 497; tuberosa, 499; ventricosa, 498
Iris, 273
Isocoma **Wrightii,** 152
Isoetes, 189; flaccida, 485; Gravesii, 204; Harveyi, 205; heterospora, 204; hieroglyphica, 204; lacustris paupercula, 204; occidentalis, 204; paupercula, 204; Tuckermani, 205
Ivy, Influence of moisture upon the formation of roots by cuttings of, 93

Jacquemontia, 313; abutiloides, 315; acrocephala, 313; azurea, 313; bracteosa, 313; Choisyana, 313; eriocephala, 314; eriocephala maynensis, 314; lactescens, 314; montana, 314; pycnocephala, 314; rufo-velutina, 314; serrata, 314; **simulata,** 314; sphaerocephala, 314; sphaerostigma, 313; tamnifolia, 314
Jeffersonia diphylla, 531
Jubula pennsylvanica, 132
Juglans, 170, 328; arctica, 170
Juncoides pilosum, 527
Juncus, 233, 515; aristulatus, 232; aristulatus pinetorum, 232; biflorus, 232; brachycarpus, 233; dichotomus, 233; marginatus, 232; marginatus biflorus, 232; marginatus pinetorum, 232; megacephalus, 233; scirpoides, 233; scirpoides **compositus,** 233; scirpoides macrostemon, 233
Jungermannia lineata, 17
Juniperus, 168; hypnoides, 168; macilenta, 168

Kantia Trichomanis, 131
KERN, F. D. & ARTHUR, J. C. North American species of Peridermium, 403
KIRKWOOD, J. E. The pollen-tube in some of the Cucurbitaceae, 327
KRAEMER, H. Studies on color in plants, 77
Kuhneola, 518

Lagenaria, 330
Larix, 405, 412, 520; decidua, 412, 437
Lathyrus dissitifolius, 144; **incanus,** 144; ornatus incanus, 144
Laurophyllum angustifolium, 178
Laurus Hollickii, 178; plutonia, 178
Ledum, 429
Lejeunea, 2, 8, 9; aneogyna, 2; Berteroana, 12, 13, 24; confluens, 2; contigua, 19; decidua, 6, 24; emarginuliflora, 14, 24; glaucescens, 131; lineata, 16, 17, 24; macroloba, 19, 21; macroloba laxior, 21; pallescens, 17; paucispina, 134; phyllobola, 15, 24; Schwaneckii, 21; versifolia, 12, 24
Lentinus cyathiformis, 215; lepideus, 215; **microspermus,** 216; **obconicus,** 215
Lepidium **crenatum,** 141
Lepidotis, 102, 113; convoluta, 116, 117
Lepiota arenicola, 213; cristata, 213; **nudipes,** 213
Leptilon, 154
Leptobryum pyriforme, 220
Leptochilus, 603, 604; contaminoides, 605; serratifolius, 604
Leptodactylon **Nuttallii,** 149; **Watsoni,** 149
Leptosphaeria **Lythri,** 220
Lesquerella Fendleri, 142; **stenophylla,** 142
Leucelene ericoides serotina, 153; **hirtella,** 153; **serotina,** 153
Lewisia, 140
Libocedrus, 166
Ligusticum montanum tenuifolium, 148
Lilium, 273, 275–277, 283; canadense, 297, 298; lancifolium, 298; longiflorum, 299; Martagon, 302; philadelphicum, 302; tigrinum, 302
Limnia, 138, 139; **arenicola,** 138;

asarifolia, 138; **depressa,** 139; **parviflora,** 139; perfoliata, 139; sibirica, 138
Limnocharis, 328; emarginata, 339
Lindsaya, 483
Linosyris serrulata, 152; viscidiflora latifolia, 152; Wrightii, 152
Liriodendron, 174; **morganensis,** 174
Literature, American botanical (1901–1903), 207; (1905), 59, 263, 447; (1906), 125, 319, 353, 397, 505, 541, 581, 617
Lithophyllum africanum, 579, 580; **Antillarum,** 579, 580; craspedium, 579, 580; daedaleum pseudodentatum, 578; platyphyllum, 578
Lithospermum carolinense, 241, 242; carolinianum, 241, 242; Gmelini, 241; hirtum, 241
Litsea falcifolia, 180
LLOYD, F. E. & UNDERWOOD, L. M. The species of Lycopodium of the American tropics, 101
Localities visited by Charles Wright, Southwestern, 561
Lomaria, 36, 592; aculeata, 40; **fraxinea,** 49; gracilis, 41; Haenkeana, 38; integrifolia, 49; juglandifolia, 38; longifolia, 197, 602; marginata, 595; **Meyeriana,** 39; procera, 592; scandens, 38; spectabilis, 47; tenuifolia, 39; variabilis, 50
Lomariobotrys, 39; Meyeriana, 39; tenuifolia, 39
Lomariopsis, 35–37, 42, 45, 591; Boryana, 47; Brackenridgei, 45; cochinchinensis, 46; cuspidata, 50; Fendleri, 595; guatemalensis, 600; guineensis, 46; Hügelii, 46; leptocarpa, 47; ludens, 40; Novae-Caledoniae, 49; palustris, 39; Pervillei, 49; phlebodes, 598; Prieuriana, 599; recurvata, 600; Seemannii, 49; Smithii, 50; sorbifolia resectum, 46; spectabilis, 47; spinescens, 40; variabilis, 50; vestita, 600; Wrightii, 601
Lonchitis repens, 192
Lonicera involucrata, 152
Lotus, 144; huministratus, 144
Luffa maxima, 331
Lunularia cruciata, 130
Luzula saltuensis, 527, 536
Lycium Berlandieri, 368
Lycogala, 557
Lycopodium: the species of the American tropics, 101
Lycopodium, 101–124, 329; acerosum, 108; adpressum, 484; alopecuroides, 229, 484; aqualupianum, 114; arboreum, 117; bifidum, 104; callitrichaefolium, 114; capillaceum, 116; carolinianum, 120–122, 484; cernuum, 116, 117, 484; clavatum, 119, 122; compactum, 102; contiguum, 120; crassum, 102; **cuernavacense,** 103, 110; cuneifolium, 114; curvatum, 117, 118; **densifolium,** 102, 106; dichaeoides, 114, 115; dichotomum, 103, 107, 111, 112; Eichleri, 119; erythraeum, 102; eversum, 104; Fawcettii, 122; filiforme, 102; funiforme, 103, 111; Gayanum, 122, 123; **goyazense,** 121; grenadense, 111; guadalupianum, 114; Hartwegianum, 102; Heeschii, 116; hippurideum, 103, 107; **Holtoni,** 122, 123; Hupeanum, 114; intermedium, 105; **iuliforme,** 120; **Jenmani,** 103, 112; Jussiaei, 122, 123; Lechleri, 107; lindsaeaceum, 122, 123; linifolium, 103, 112, 113; lucidulum, 102; marianum, 116; **meridionale,** 121; **montanum,** 103, 107, 111; Moritzii, 114; nitens, 103, 109, 110; **Orizabae,** 103, 110; passerinoides, 109; pendulinum, 117, 119; pendulum, 119; Picardae, 103, 110; pinnatum, 229; pithyoides, 103, 109; **polycarpum,** 102, 105; porophilum, 102; **portoricense,** 103, 108; **Pringlei,** 103, 109; **prostratum,** 229; reflexum, 102, 104–107; reflexum densifolium, 106, 107; reflexum polycarpum, 105; reversum, 104; rigidum, 103–106; **roraimense,** 114, 115; rufescens, 102; scariosum, 122, 123; secundum, 116; Selago, 102; setaceum, 103, 108, 112; Sieberianum, 102, 105; sikkimense, 114; squarrosum, 104; subulatum, 114, 115; subulatum pastoense, 115; taxifolium, 103, 109, 110; **tenuicaule,** 103, 113; tetragonum, 102; thyoides, 122; **tortum,** 117, 118; verticillatum, 108, 109; vulcanicum, 116; **Williamsii,** 103, 112; **Wilsonii,** 103, 111
Lygodesmia, 161; exigua, 161
Lygodium japonicum, 191
Lythrum alatum, 221

Macbridea, 244; pulchella, 244; pulchra, 244
MACKENZIE, K. K. Notes on Carex, 439
Macrolejeunea, 16, 17; lineata, 16, 17; pallescens 17
Madronella **dentata,** 150; **parvifolia,** 150
Magnolia, 170; auriculata, 174; glauca, 530; pyramidata, 240; tenuifolia, 174, 182
Mahonia, 140; Aquifolium nutkana, 141

Malapoenna falcifolia, 180
Malus, 310
Malva, 490
Malvastrum elatum, 311; **exhumatum,** 311
Mamillaria aggregata, 146; missouriensis caespitosa, 146; similis, 146
Marchantia polymorpha, 130, 132
Marsonia Potentillae **Helleri,** 219
MATHEWSON, C. A. The behavior of the pollen-tube in Houstonia coerulea, 487
Mechanism of seed-dispersal in Polygonum virginianum, 377
Melampsora, 520; **albertensis,** 517
Melampsorella, 405, 407, 408; Cerastii, 405, 435
Melampsoridium, 405–407; betulinum, 437
Melampsoropsis, 406
Melissa caroliniana, 243; grandiflora, 244
Melittis caroliniana, 243, 244
Melo, 330
Melothria, 327, 331–336, 338, 342; pendula, 331, 338
Meniscium, 198, 469; cuspidatum, 471; reticulatum, 198, 469
Mentha arvensis Penardi, 150; **Penardi,** 150
Merremia, 495; dissecta, 500
Mertensia **platensis,** 150; polyphylla platensis, 150
Mesadenia sulcata, 536
Mesotaenium, 346, 349–351, 607; Braunii, 346; chlamydosporum, 346, 351; chlamydosporum minor, 346; DeGreyi, 349, 351, 607; DeGreyi breve, 346; Endlicherianum, 347, 351, 607; macrococcum, 346, 351; macrococcum micrococcum, 346; micrococcum, 346; **minimum,** 347, 351
Mesozoic flora of the Atlantic coastal plain, 163
Metzgeria conjugata, 131
Micrampelis, 327, 331–336, 338, 342; lobata, 331, 338
Micrasterias, 611; conferta Novaeterrae, 611; denticulata, 611; denticulata angulosa, 611; pinnatifida, 611; truncata, 611
Moisture, Absorption of atmospheric, 367
Moisture: its influence upon the formation of roots by cuttings of ivy, 93
Momordica, 330
Monardella dentata, 150; parvifolia, 150
Monilia **Avenae,** 219
Montia, 138, 139, fontana, 138
Montiastrum, 138
Moriconia cyclotoxon, 165
Mucilago, 557
Mucor, 557–559; ascophorus, 559, 560; caninus, 560; Mucedo, 557–560; sphaerocephalus, 559, 560; stercoreus, 560; stolonifer, 559, 560; vulgaris, 558–560
Mucor Mucedo, The identity of, 557
Muhlenbergia, 230; trichopodes, 231
Myrica, 308; alkalina, 309; carolinensis, 528, 530, 533; cerifera, 308; Gale, 30, 308; **Hendersoni,** 308; longa, 170
Myzorrhiza, 151; ludoviciana, 151, v; **multiflora,** 151

Nabalus, 161
Naias major, 300
Naiocrene, 138, 139; **parvifolia,** 139
Narthecium, 278; ossifragum, 298
Nectar-glands of Vicia Faba, Cytological changes accompanying secretion in the, 247
Nelumbium arcticum, 173; Lakesianum, 174; tenuifolium, 174
Nelumbo, 173; Kempii, 173; lutea, 174; primaeva, 173
Nephrodium, 198; molle, 477
Nephrolepis biserrata, 483; exaltata, 484
Netrium, 348, 350, 351, 608; Digitus, 348, 351, 608; interruptum, 350, 351; Naegelii, 349, 351, 608; oblongum, 349, 351; oblongum cylindricum, 349; oblongum f. major 349
Neviusia alabamensis, 532
New coralline algae from Culebra, Two, 577
New England desmids of the subfamily Saccodermae, 343
New or otherwise noteworthy plants from the coastal plain of Georgia, 229
New southwestern species of Pentstemon, Two, 445
New species of fungi, 213
New species of Uredineae, 27, 513
Newfoundland, Some desmids from, 607
North American Convolvulaceae, Studies in the, 313, 495
North American Polygonaceae, Studies in, 51
North American species of Calceolaria, 547
North American species of Peridermium, 403
Notes on Calochortus, 537
Notes on Carex, 439
Notes on the distribution of some Alabama plants, 523
Noteworthy plants from the coastal plain of Georgia, 229

Nothaphyllon, 151
Notholaena, 374
Nymphaea advena, 235, 236; **fluviatilis,** 234, 235; macrophylla, 235; orbiculata, 236; sagittifolia, 236; variegata, 236

Odontolejeunea, 13
Odontoschisma prostratum, 129, 131
Odontosoria, 483; clavata, 200, 458, 483
Odostemon, 140; **Aquifolium,** 141; **Fremontii,** 141; **nervosus,** 141; **nutkanus,** 141
Oenothera alyssoides minutiflora, 146; idahoensis, 146; marginata, 146; pallida latifolia, 146; strigulosa pubens, 146
Olfersia, 603; cervina, 603; corcovadensis, 603; Kunzeana, 196, 197, 598
Oligoneuron, 152
Onoclea, 295, 302; scandens, 38
Operculina, 495, 500, 501; **aegyptia,** 497, 502; alata, 496–498; **alatipes,** 496, 498, 499; altissima, 497; **ampliata,** 503; **angustiloba,** 497, 501; **codonantha,** 503; Convolvulus, 496, 499; **coptica,** 500; **dissecta,** 497, 500; leptoptera, 496, 498; macrocarpa, 495, 496, 499; **Palmeri,** 497, 502; Pavoni, 503; **platyphylla,** 497, 502; populifolia, 497; pterodes, 497; **rhodocalyx,** 496, 498, 499; **Roseana,** 497, 500–502; **rubicunda,** 496, 498; triquetra, 496, 497; tuberosa, 497, 499; ventricosa, 496, 498
Operculina, The genus, 495
Ophioglossum, 459; japonicum, 191; palmatum, 459; pusillum, 459
Oreobroma, 140
Oreocarya affinis perennis, 150; **hispidissima,** 150; **perennis,** 150
Oreochrysum, 152; **Parryi,** 153
Orobanche, 151; ludoviciana, 151; multiflora, 151
Osmanthus americanus, 535
Osmunda, 164; cinnamomea, 462; **delawarensis,** 164, 182; palustris, 463; pollicina, 42, 49; Presliana, 164; regalis, 164, 462; trifrons, 50
Osmundites, 164; skidegatensis, 164
Otiona Aitonia, 129
Oxytropis Hallii, 144

Pachylophus **marginatus,** 146
Paliurus **populiferus,** 177
Pallavicinia Lyellii, 131
Palmia, 318
Palmogloea macrococca, 346; micrococca, 346
Panaeolus, 218
Parkinsonia, 374; microphylla, 368, 374
Parnassia asarifolia, 532
Paronychia **brevicuspis,** 137; riparia, 530; sessiliflora brevicuspis, 137
Parthenium, 156
PECK, C. H. New species of fungi, 213
Pecopteris kudlisetensis, 166
Pedicularis, 274
Pedunculate species of Trillium, The, 387
Pellaea andromedaefolia, 30; atropurpurea cristata, 190; gracilis, 30
Penium, 608; Brebissonii, 347; crassum, 347; Digitus, 348; interruptum, 350; lamellosum, 348; Libellula intermedium, 608; Naegelii, 349; oblongum, 349
Pentstemon, 445, 446; acuminatus, 150, 151; angustifolius, 151; angustifolius **caudatus,** 151; antirrhinoides, 446; barbatus, 151; barbatus trichander, 151; caudatus, 151; cyathophorus, 151; Fendleri, 151; **jacintensis,** 445; **Plummerae,** 445, 446; Rothrockii, 445; secundiflorus, 150, 151; Torreyi, 151; **trichander,** 151; **unilateralis,** 150
Pentstemon, Two new southwestern species of, 445
Pepo, 330
Peridermium, 403–438, 514, 518, 520; abietinum, 409, 411, 429, 430; abietinum decolorans, 428; acicolum, 409, 410, 413; balsameum, 410, 412, 435; **boreale,** 409, 411, 425; carneum, 409, 410, 414, 416; Cerebrum, 405, 409, 411, 423, 425, 514; **coloradense,** 409, 411, 426, 427; columnare, 406, 409, 412, 432; conorum, 431; **conorum-Piceae,** 409, 411, 431; **consimile,** 409, 411, 427; decolorans, 409, 411, 427–430; deformans, 423; **delicatulum,** 408, 410, 412; elatinum, 405, 410, 412, 434; Engelmanni, 431; Ephedrae, 410, 412, 437; filamentosum, 405, 409, 410, 418, 419; **fusiforme,** 409, 411, 421; giganteum, 423; **globosum,** 409, 411, 424; **gracile,** 409, 410, 417; Harknessii, 409, 410, 421; Holwayi, 409, 411, 431, 514; **intermedium,** 409, 410, 416; **Laricis,** 410, 412, 436; **mexicanum,** 409, 411, 422; **montanum,** 408, 410, 413; oblongisporium Ravenelii, 414; orientale, 416; ornamentale, 409, 412, 433; Peckii, 410, 412, 433, 518; Pini, 405; Pini minor, 437; **pseudo-balsameum,** 409, 412, 430, 514; pyriforme, 409, 410, 419; Ravenelii, 414; Rostrupi, 409, 410, 416; **stalactiforme,** 409, 410, 419; Strobi, 425

Peridermium, North American species of, 403
Peritoma **Sonorae,** 142
Persea, 67, 69; gratissima, 67, 76
Persicaria **omissa,** 57
Persimmons, Tannin cells of, 567
Persoonia Lesquereuxii, 173
Phacelia **ciliosa,** 149; sericea ciliosa, 149
Phaseolites manhassettensis, 176
Phaseolus, 32
Phegopteris, 30, 476; reptans, 477
Phelipaea, 151
Phileozera multiflora, 157
Phlebodium aureum, 467
Phlox, 221; **depressa,** 149; multiflora depressa, 149
Phoenix dactylifera, 261
Phryma caroliniensis, 242
Phymatodes, 603; exiguum, 458, 466
Physopella, 520
Phytolacca, 377
Picea, 169, 405, 411; alba, 432; canadensis, 411, 429, 432; **cliffwoodensis,** 169; Engelmanni, 411, 425, 426, 428, 432; excelsa, 169, 411, 430, 432; Mariana, 411, 426–428, 432; Parryana, 411, 425; pungens, 425; rubens, 427, 428, 432; rubra, 411, 427, 428, 432; sitchensis, 411, 429
Picradenia, 156; macrantha, 156; multiflora, 157; odorata Osterhoutii, 157; pumila, 156
Pieris phillyreifolia, 534
Pinites, 169
Pinus, 169, 295, 299, 329, 405–407, 410, 413, 422, 520; australis, 415; Banksiana, 423; contorta, 410, 421; divaricata, 411, 423; echinata, 410, 411, 416, 424; Elliottii, 410, 415, 457; filifolia, 410, 418; heterophylla, 415; inops, 420; insignis, 410, 421; Jeffreyi, 410, 419; Laricio, 299; mitis, 416, 424; Murrayana, 410, 413, 419, 421; oocarpa, 411, 423; palustris, 410, 411, 415, 416, 422, 519, 529, 533; patula, 411, 423; ponderosa, 410, 411, 418, 421, 424, 514; rigida, 410, 411, 414, 417, 420, 423; scopulorum, 410, 413; serotina, 415, 524; Strobus, 424; sylvestris, 298, 410, 420; Taeda, 410, 411, 413, 415, 422, 424, 519; virginiana, 410, 411, 420, 423, 524
PIPER, C. V. Notes on Calochortus, 537
Piptostegia, 496
Plagiochila Smallii, 131
Planera aquatica, 534; Ungeri, 310
Plants, Studies on color in, 77
Plants from the coastal plain of Georgia, 229
Plants from Florissant, Fossil, 307
Platanus Kümmelii, 174
Platyschkuhria, 154; **integrifolia,** 155; **oblongifolia,** 155
Pleospora **magnifica,** 221
Pleuroschisma prostratum, 129
Pleurotaenium, 609; coronatum, 609; Ehrenbergii, 609
Pluchea, 154; borealis, 154; purpurascens, 517
Podophyllum, 276
Podosemum, 230
Poeciloptcris, 604; crenata, 603–605
Pogonatum alpinum, 219
Pogonia ophioglossoides, 462
Poikilopteris, 504, 50, 604, 605
Pollen-tube in Houstonia coerulea, The behavior of the, 487
Pollen-tube in some of the Cucurbitaceae, 327
Polybotrya, 603; articulata, 40
Polygala nana, 533
Polygonaceae, Studies in North American, 51
Polygonella, 234; gracilis, 234
Polygonum, 377; alpinum, 379; Bistorta oblongifolium, 57; bistortoides, 57; **buxiforme,** 56; littorale, 56; omissum, 57; ramossissimum, 57; **rubescens,** 56; scandens, 379; virginianum, 377–386
Polygonum virginianum, The mechanism of seed-dispersal in, 377
Polypappus, 154; sericeus, 154
Polypodium, 44; amplum, 198; angustifolium, 192; binerve, 44; costatum, 192; hesperium, 191; orbiculatum, 45; palustre, 38; pectinatum, 465, 467; Plumula, 466, 467; polypodioides, 458, 466; reticulatum, 198; vulgare, 191
Polystichum acrostichoides, 472
Populus, 172, 517; acerifolia, 312; aequalis, 312; apiculata, 172; balsamoides latifolia, 307; cordifolia, 312; **dakotana,** 312; Heerii, 307; laevigata, 312; **Lesquereuxii,** 307; **Newberryi,** 312; **Scudderi,** 307; tremuloides, 517
Poronia **macrospora,** 220
Portulaca oleracea, 329
Potentilla, 143; anserina concolor, 143; bipinnatifida, 143; bipinnatifida platyloba, 143; ovalis, 143; **platyloba,** 143; prostrata, 143; rubricaulis, 143; **rubripes,** 143
Pothos, 328
Prenanthella, 160; **exigua,** 161
Prenanthes, 161; exigua, 161
Prionolejeunea, 3, 133
Proserpinaca, 238; palustris, 239; pectinata, 239; platycarpa, 239
Prosopis, 374

Prunus Copalin, 144; demissa, 143; **melanocarpa,** 143, 144; virginiana, 144
Psathyrella, 218; **angusticeps,** 217
Pseudocymopterus montanus multifidus, 147; montanus purpureus, 147; **multifidus,** 147; **purpureus,** 147; **tenuifolius,** 147
Pseudotsuga, 405, 411, 514; Douglasii, 431; mucronata, 411, 431, 514
Psilotum nudum, 485
Pteridium, 251, 252, 259; aquilinum, 261
Pteridophytes observed during three excursions into southern Florida, 455
Pteris, 272; aquilina pseudocaudata, 471; caudata, 472; longifolia, 457, 461, 472
Puccinia, 513, 522; Agropyri, 29; atra, 519; Cirsii, 33; Cnici, 33; **Dolichi,** 28, 513; Echinopteridis, 516; Eleocharidis, 28, 29; **Fimbristylidis,** 28; **inflata,** 516; **Pattersoniana,** 29; suaveolens, 33; **Tridacis,** 516; Xanthii, 517
Pucciniastrum, 405–407, 518
Puerto Rico, Hepaticae of, 1
Pycnolejeunea, 1, 2, 4, 9, 19; macroloba, 19, 23, 24; Schwaneckei, 21–25; Spruceana, 19
Pyrola, 432

Quercus, 309, 328, 329, 339, 424; coccinea, 311; Fendleri, 309; laurifolia, 529; Phellos, 529; **Ramaleyi,** 309

Radula pallens, 129, 131; pycnolejeuneoides, 22
Randia, 521, 522
RANDOLPH, H. The influence of moisture upon the formation of roots by cuttings of ivy, 93
Reboulia hemisphaerica, 129, 130, 132
Rectolejeunea, 1, 4, 8, 9, 19, 20; **Berteroana,** 12–15; **emarginuliflora,** 14, 15, 25; **flagelliformis,** 9, 10, 13–15, 25; **phyllobola,** 15, 131
Reduction, A study of synapsis and, 271
REED, H. S. & SMOOT, I. The mechanism of seed-dispersal in Polygonum virginianum, 377
Rhamnus ellipticus, 311; **Kirchneri,** 311
Rhapidophyllum, 526; Hystrix, 526
Rhexia Alifanus, 238; glabella, 238
Rhizopus, 559; nigricans, 558–560
Rhododendron, 534; Cuthbertii, 240, 534; punctatum, 534
Rhynchospora, see Rynchospora
Riccardia latifrons, 129, 130; multifida, 131; palmata, 129
Richardsonia, 487, 489; pilosa, 487
Rocky mountain flora, Studies on the, 137
Rumex occidentalis, 137; **praecox,** 137
Kussula **nigrescentipes,** 214; ochrophylla, 215; **subvelutina,** 215; uncialis, 215
RYDBERG, P. A. Studies on the Rocky mountain flora, 137
Rydbergia **Brandegei,** 156; glabrata, 156
Rynchosia texana, 27
Rynchospora leptorhyncha, 231; semiplumosa, 526

Saccodermae, New England desmids of the sub-family, 343
Sagenia, 200
Salix, 308; amygdalaefolia, 307; cordata Watsonii, 137; flava, 137; flexuosa, 171; myrtilloides, 308; proteaefolia, 171, 182; proteaefolia flexuosa, 171; proteaefolia longifolia, 171; **Ramaleyi,** 307; **Watsonii,** 137
Salomonia, 274, 289; biflora, 286, 305
Sambucus, 308, 377
Sanilum, 318
Santalum, 182; americanum, 309; cognatum, 182; ellipticum, 182; **Novae-Caesareae,** 182
Sapotacites, 181; **Knowltoni,** 181, 182; obovata, 181; retusa, 181
Sarcoscypha **dawsonensis,** 220
Sarracenia, 236, 237; flava, 237; flava × minor, 236, 237; formosa, 237; minor, 236, 237; **minor × psittacina,** 236; psittacina, 236, 237
Schizaea, 191, 462; Germani, 461; pusilla, 191
Schkuhria, 154; integrifolia, 155
Scirpus atrovirens, 526; carinatus, 525; georgianus, 526
Secretion in the nectar-glands of Vicia Faba, Cytological changes accompanying, 247
Seed-dispersal in Polygonum virginianum, The mechanism of, 377
Selaginella, 189, 190, 204; acanthonota, 202; arenicola, 485; Bolanderi, 203; Bourgeaui, 203; caribensis, 204, 485; densa, 204; Engelmanni, 204; Fendleri, 203; Hanseni, 202; Haydeni, 203; montanensis, 203; **Parishii,** 202; rhodospora, 485; rupestris, 202; rupestris Fendleri, 203; Schmidtii, 203; Schmidtii Krauseorum, 203; Sherwoodi, 202; Underwoodii, 203; Wallacei, 203; Wrightii, 203
Selago, 102
SELBY, A. D. Studies in etiolation, 67
Senecio aureus, 159; canus, 158; cro-

catus, 159; cymbalarioides, 159; Douglasii, 160; exaltatus, 158; fedifolius, 159; **Flintii,** 157; **Harbourii,** 158; integerrimus, 158; **multicapitatus,** 160; **oodes,** 158; pentodontus, 160; Purshianus, 158; pyrrochrous, 159; Riddellii, 160; salignus, 520; **Tracyi,** 159; **turbinatus,** 159
Sequoia gracillima, 165, 168; heterophylla, 165; Reichenbachi, 165, 168
Shrubs, Absorption of atmospheric moisture by desert, 367
Shutereia, 318; bicolor, 318; **sublobata,** 318
Shuteria, 318
Sida lepidota sagittaefolia, 145; **sagittaefolia,** 145
Siphonychia pauciflora, 530
SMALL, J. K. Studies in North American Polygonaceae, 51
SMOOT, I. & REED, H. S. The mechanism of seed-dispersal in Polygonum virginianum, 377
Solanum rostratum, 150
Solea, 547, 548; glutinosa, 549; longifolia, 554; riparia, 554; stricta, 553; verbenacea, 553; verticillata, 547, 549
Solidago, 152; canadensis gilvocanescens, 153; canadensis scabra, 153; canadensis scabriuscula, 153; **gilvocanescens,** 153; **pallida,** 153; Parryi, 153; scabra, 153; **scabriuscula,** 153; speciosa pallida, 153
Some desmids from Newfoundland, 607
Some new or otherwise noteworthy plants from the coastal plain of Georgia, 229
Sorbus occidentalis, 521
Southern Florida, Pteridophytes observed during three excursions into, 455
Southwestern localities visited by Charles Wright, 561
Southwestern species of Pentstemon, Two new, 445
SPALDING, V. M. Absorption of atmospheric moisture by desert shrubs, 367
Spartina, 231; Bakeri, 231; juncifórmis, 231
Species of Calceolaria, North American, 547
Species of Lycopodium of the American tropics, The, 101
Species of Pentstemon, Two new southwestern, 445
Species of Peridermium, North American, 403
Species of Stenochlaena, American, 591; Old-world, 35
Species of Trillium, The pedunculate, 387
Sphaeralcea incana, 145; **marginata,** 145; Munroana, 145
Sphaeropsis, 219
Sphaerostigma **minutiflora,** 146; **pubens,** 146
Sphaerozosma, 615; excavatum, 615
Sphagnum, 118
Spiranthera, 496, 500; pentaphylla, 502
Spirotaenia, 343, 345, 350; condensata, 345, 350; obscura, 350; obscura **crassa,** 345; parvula, 345, 350
Sporobolus, 230; Curtissii, 230; floridanus, 230; **teretifolius,** 229, 230, 237
Staphylea trifolia, 533
Status of Poecilopteris crenata Presl, The, 603
Staurastrum, 614; Arctiscon, 615; arcuatum, 614; aristiferum, 614; crenulatum, 614; dubium, 614; echinatum, 614; eustephanum, 615; gracile, 614; Johnsonii, 614; longispinum bidentatum, 614; macrocerum, 614; nanum, 614
Stegania, 36
Stellaria, 435
Stenochlaena, 35–50, 197, 464, 591, 593, 595, 602, 603; aculeata, 40, 41; **angusta,** 594, 596; Blumeana, 38; **Brackenridgei,** 43–45; **buxifolia,** 42, 45; **cochinchinensis,** 45, 46; **decrescens,** 42, 43, 46; **erythrodes,** 593, 595, 598, 603; feegeensis, 45; **Fendleri,** 593, 595, 597, 601; gracilis, 40, 41; **guineensis,** 43, 44, 46; Haenkeana, 38; **Hugelii,** 44, 46, 48; **jamaicensis,** 593, 595, 597; japurensis, 593, 598, 599; **Kunzeana,** 196, 465, 592, 593, 596, 598; laurifolia, 37, 38; **leptocarpa,** 44, 47; **lomarioides,** 42, 43, 47; **Mannii,** 42, 43, 47; marginata, 595; **Maxoni,** 594, 597, 599, 603; Meyeriana, 39; **Milnei,** 37, 38; **Novae-Calendoniae,** 44, 48, 49; oleandrifolia, 45, 48, 49; palustris, 37, 38; **Pervillei,** 43, 44, 49; Pittieri, 600; **pollicina,** 42, 48, 49; **Prieuriana,** 593, 599; **recurvata,** 594, 600; scandens, 38; **Seemannii,** 45, 49; **Smithii,** 45, 47, 50; sorbifolia, 43, 44, 46, 48, 592, 594, 597, 600, 602; **tenuifolia,** 39; **variabilis,** 44, 50; **vestita,** 594, 597, 600; **Williamsii,** 40, 41; Wrightii, 593, 596, 601
Stenotus, 152
Sterculia, 178; cliffwoodensis, 178; lugubris, 178; **minima,** 177; mucronata, 177, 178

Stigmatophyllon periplocifolium, 516
STOCKARD, C. R. Cytological changes accompanying secretion in the nectar-glands of Vicia Faba, 247
Streptanthus, 142; cordatus, 142; crassifolius, 142; longirostris, 142
Strobus, 411; Strobus, 411, 424
Strophostyles, 32
Strumpfia, 223, 227
Struthiopteris, 197
Studies in etiolation, 67
Studies in North American Convolvulaceae, 313, 495
Studies in North American Polygonaceae, 51
Studies on color in plants, 77
Studies on the Rocky Mountain flora, 137
Study of synapsis and reduction, A, 271
Stylodon, 243; scabrum, 243
Svida, 147
Synapsis and reduction, A study of, 271

Tannin cells of persimmons, 567
Taxilejeunea, 17
Taxodium distichum, 525, 528, 534; imbricarium, 237
Tectaria, 200, 603; **Amesiana,** 479; **coriandrifolia,** 199, 200, 480; **heracleifolia,** 199, 200, 481; **minima,** 199, 480, 482; trifoliata, 199, 200, 482
Telaranea nematodes, 130, 131
Teratophyllum, 37, 40; aculeatum, 40, 41; articulatum, 40
Tessaria, 154; borealis, 154
Tetmemorus, 609; Brebissonii minimus, 609; Brebissonii minor, 609; Brebissonii turgidus, 609; granulatus, 609; laevis, 609; minutus, 609
Tetragonolobus, 144
Tetraneuris, 155, 156; argentea, 155; fastigiata, 155; glabra, 155; **glabriuscula,** 155; herbacea, 155; Ivesiana, 155; leptoclada, 155 stenophylla, 155, v
Thaspium barbinode Chapmani, 240; montanum tenuifolium, 147
Thelypodium crenatum, 141; integrifolium, 141
Thladiantha, 332
Three excursions into southern Florida, Pteridophytes observed during, 455
Thuja, 166; cretacea, 169
Thujopsis, 166
Thyella, 313; **acrocephala,** 313; **bracteosa,** 313; **Choisyana,** 313; **eriocephala,** 314; **lactescens,** 314; **maynensis,** 314; **montana,** 314; **pycnocephala,** 314; **rufovelutina,** 314; **serrata,** 314; **sphaerocephala,** 314; tamnifolia, 313, 314
Thymbra, 244; caroliniana, 244
Thymus carolinianus, 243, 244; grandiflorus, 243
Tilia, 216
Tillandsia usneoides, 527
Tithymalus arkansanus **coloradensis,** 145
Todites, 164
Tovara, 378
Trachelospermum difforme, 535
Trachylejeunea, 16, 17
Tradescantia, 336
Trichochloa, 230
Trichomanes, 201; Krausii, 201, 460; punctatum, 461; reptans, 461; sphenoides, 201, 460
Tridax procumbens, 517
Trillidium Govanianum, 393
Trillium, 273, 387–396; acuminatum, 396; affine, 389, 396; album, 392; brevipetalum, 396; californicum, 395; camschatcense, 389, 393; camtschaticum, 393; Catesbaei, 387, 389, 390, 395, 396; cernuum, 387, 388, 390, 392, 395, 396; crassifolium, 395; **declinatum,** 388–390, 392, 394; divaricatum, 396; erectum, 387, 388, 390–393, 396; erectum album, 392, 393; erectum atropurpureum, 392; erectum declinatum, 389; erectum flavum, 392; erectum japonicum, 393; erectum viridiflorum, 392; erythrocarpum, 394; flavum, 396; foetidum, 392; Govanianum, 389, 393; grandiflorum, 387, 389, 392, 394, 395; hamosum, 396; latifolium, 396; lirioides, 396; nervosum, 395; nivale, 387–389, 393, 394; nutans, 396; obcordatum, 396; obovatum, 392–394; ovatum, 389, 395; pendulum, 391, 392; pictum, 394; pumilum, 395; purpureum, 392; pusillum, 389, 395; rhomboideum, 391; rhomboideum album, 391; rhomboideum grandiflorum, 394; rivale, 387, 389, 393; Rugelii, 388, 391; **Scouleri,** 389, 394, 395; **simile,** 388, 391; Smallii, 389, 392; spatulatum, 396; stylosum, 395; texanum, 395; Tschonoskii, 389, 392, 393; undulatum, 388, 389, 393, 394; Vaseyi, 388, 390
Trillium, The pedunculate species of, 387
Triosteum angustifolium, 33
Tropics, The species of Lycopodium of the American, 101
Tsuga, 295, 405, 412, 524; canadensis, 301, 412, 434, 518, 524, 525; Mertensiana, 518
Tuberculariа carnea, 414

Tubiflora, 242
Turpithum, 496
Two new coralline algae from Culebra, 577
Two new southwestern species of Pentstemon, 445
Typha latissima, 307; **Lesquereuxii,** 307

Ulmus, 492; fulva, 219
UNDERWOOD, L. M. American ferns, 189, 591; The genus Stenochlaena, 35
UNDERWOOD, L. M. & LLOYD, F. E. The species of Lycopodium of the American tropics, 101
Unifolium, 274
Uredineae, New species of, 27, 513
Uredinopsis, 32
Uredo **biocellata,** 517; **Chaetochloae,** 518; **Dichromenae,** 31; **Dolichi,** 513; **Holwayi,** 518; ledicola, 429; superior, 29
Uromyces, 28, 513, 522; appendiculatus, 32; Clitoriae, 515; **Dolicholi,** 27; **Eleocharidis,** 514; **insularis,** 515; Junci, 515
Utricularia, 607
Uvularia floridana, 528

Vaccinium, 406, 432; myrtilloides, 148; Myrtillus, 148; **oreophilum,** 148
Verbena, 243, 244; carnea, 242, 243; Carolina, 242, 243; carolinensis, 242, 243; caroliniana, 242, 243
Verbesina encelioides exauriculata, 154
Vernonia, 183–188; **arctata,** 184–187; bahamensis, 183–188; Blodgettii, 185; cinerea, 183, 184; **insularis,** 184, 185; longifolia, 183; longifolia Sintenisii, 183; longifolia Vahliana, 183; **obcordata,** 184, 185, 187, 188
Vernonia in the Bahamas, 183
Vesicaria stenophylla, 142
Vicia **dissitifolia,** 144; Faba, 247–262
Vigna, 32
Viola, 547, 548; Calceolaria, 548; glutinosa, 549; lutea, 84; polygalaefolia, 549; stricta, 553; verticillata, 547, 549
Vittaria lineata, 466

Widdringtonites, 168; Reichii, 167, 169
WILSON, G. W. The identity of Mucor Mucedo, 557
Woodwardia, 523; areolata, 472; virginica, 473
WOOTON, E. O. Southwestern localities visited by Charles Wright, 561
Wright, Southwestern localities visited by, 561

Xanthidium, 613; antilopaeum, 613; antilopaeum minneapoliense, 613; armatum, 613; cristatum, 613
Ximenesia **exauriculata,** 154
Xylosteum involucratum, 152

Zea Mays, 261



1-9. CHEILOLEJEUNEA DECIDUA (Spruce) Evans

10-25. RECTOLEJEUNEA FLAGELLIFORMIS Evans



1-8. RECTOLEJEUNEA EMARGINULIFLORA (Gottsche) Evans

9-23. PYCNOLEJEUNEA SCHWANECKEI (Steph.) Schiffn.



CYSTOLEJEUNEA LINEATA (Lehm. & Lindenb.) Evans



STUDIES IN ETIOLATION



STUDIES IN ETIOLATION



CROSSOTOLEJEUNEA BERMUDIANA Evans

MAGNOLIA, SALIX, CINNAMOMUM, FICUS



SAPOTACITES, OSMUNDA, COLUTEA

BRACHYPHYLLUM



NECTAR-GLANDS OF VICIA FABA



NECTAR-GLANDS OF VICIA FABA



SYNAPSIS AND REDUCTION IN ACER



SYNAPSIS AND REDUCTION IN ACER AND SALOMONIA



SYNAPSIS AND REDUCTION IN SALOMONIA AND GINKGO



SYNAPSIS AND REDUCTION IN GINKGO AND BOTRYCHIUM



1,2,7. MELOTHRIA. 3,4,8. MICRAMPELIS. 5,6. CYCLANTHERA



9



10



11



12



13





9,10. MELOTHRIA. 11. MICRAMPELIS. 12-15. CYCLANTHERA

CALCEOLARIA TENUIFOLIA Dowell.

CALCEOLARIA LONGIPES Dowell.

CALCEOLARIA GLABRA Dowell.

CALCEOLARIA BREVIS, Dowell.



CALCEOLARIA ROSEI Dowell.



GONIOLITHON ACROPETUM Fosl. & Howe.



GONIOLITHON ACROPETUM Fosl. & Howe.

LITHOPHYLLUM ANTILLARUM Fosl. & Howe.



LITHOPHYLLUM ANTILLARUM Fosl. & Howe.

GENERAL LIBRARY
UNIV. OF MICH.
FEB 9 1907

VOL. 33 DECEMBER, 1906 NO. 12

BULLETIN

OF THE

TORREY BOTANICAL CLUB

Editor

JOHN HENDLEY BARNHART

Associate Editors

ALEXANDER WILLIAM EVANS
TRACY ELLIOT HAZEN
MARSHALL AVERY HOWE
WILLIAM ALPHONSO MURRILL
HERBERT MAULE RICHARDS
ANNA MURRAY VAIL

CONTENTS

American ferns—VII. . . . . LUCIEN MARCUS UNDERWOOD. 591
Some desmids from Newfoundland . . JOSEPH AUGUSTINE CUSHMAN. 607
INDEX TO AMERICAN BOTANICAL LITERATURE 617
INDEX TO VOLUME 33 621

PUBLISHED FOR THE CLUB
THE NEW ERA PRINTING COMPANY
LANCASTER, PA.

THE TORREY BOTANICAL CLUB

President,
HENRY H. RUSBY, M.D.

Vice-Presidents,
EDWARD S. BURGESS, PH.D. LUCIEN M. UNDERWOOD, PH.D., LL.D.

Recording Secretary,
C. STUART GAGER, PH.D.
Botanical Garden, Bronx Park, New York ity.

Corresponding Secretary,
JOHN K. SMALL, PH.D.
Botanical Garden, Bronx Park, New York City

Treasurer,
CARLTON C. CURTIS, Ph.D.
Columbia University, New York City.

MEETINGS

Meetings twice each month from October to May inclusive: the second Tuesday, at 8:00 P.M., at the American Museum of Natural History; the last Wednesday, at 3:30 P.M., in the Museum Building of the New York Botanical Garden.

PUBLICATIONS

All subscriptions and other business communications relating to the publications of the Club should be addressed to the Treasurer, Carlton C. Curtis, Columbia University, New York City.

Bulletin. Monthly, established 1870. Price, $3.00 a year; single numbers 30 cents. Of former volumes, only 24–32 can be supplied separately; certain numbers of other volumes are available, but the entire stock of some numbers has been reserved for the completion of sets. Manuscripts intended for publication in the BULLETIN should be addressed to Dr. John Hendley Barnhart, Editor, New York Botanical Garden, Bronx Park, New York City.

Torreya. Monthly, established 1901. Price, $1.00 a year. Manuscripts intended for publication in TORREYA should be addressed to Dr. Marshall A. Howe, Editor, New York Botanical Garden, Bronx Park, New York City.

Memoirs. Occasional, established 1889. (See last pages of cover.)

Preliminary Catalogue of Anthophyta and Pteridophyta within 100 miles of New York City, 1888. Price, $1.00.

THE BRYOLOGIST

WITH THE JANUARY, 1906, NUMBER BEGINS ITS NINTH YEAR AND VOLUME

It is a 16–20 page bi-monthly devoted to the study of the Mosses, Hepatics and Lichens. It is fully illustrated with new, original and artistic drawings and halftones. It is indispensable to the working bryologist, professional as well as amateur. Send for sample copy. Subscription price $1.00 a year.

Address, MRS. ANNIE MORRILL SMITH, 78 Orange Street, Brooklyn, N. Y.

INDEX TO AMERICAN BOTANICAL LITERATURE

This Index, printed each month in the BULLETIN, is reissued on library catalogue cards; these are furnished to subscribers at the rate of one cent for each card.

Subscriptions to the card-issue may be addressed to the Treasurer of the Torrey Botanical Club,

CARLTON C. CURTIS, COLUMBIA UNIVERSITY, New York City.

Beginning with January, 1903, the Index has also been reprinted on paper, on one side only, and can be furnished in that form at the rate of $3.00 a year.

North American Flora

THIS work is designed to present descriptions of all plants growing, independent of cultivation, in North America, here taken to include Greenland, Central America, the Republic of Panama, and the West Indies, except Trinidad, Tobago, and Curaçao and other islands off the north coast of Venezuela, whose flora is essentially South American.

It will be published in parts at irregular intervals by the New York Botanical Garden through the aid of the income of the David Lydig Fund bequeathed by Charles P. Daly.

It is planned to issue parts as rapidly as they can be prepared, the extent of the work making it possible to commence publication at any number of points. The completed work will form a series of volumes with the following sequence:

Volume 1. Mycetozoa, Schizophyta, Diatomaceae.
Volumes 2 to 10. Fungi.
Volumes 11 to 13. Algae.
Volumes 14 and 15. Bryophyta.
Volume 16. Pteridophyta and Gymnospermae.
Volumes 17 to 19. Monocotyledones.
Volumes 20 to 30. Dicotyledones.

The preparation of the work has been referred by the Scientific Directors of the Garden to a committee consisting of Professors L. M. Underwood and N. L. Britton.

Professor George F. Atkinson of Cornell University, Professors Charles R. Barnes and John M. Coulter of the University of Chicago, Mr. Frederick V. Coville of the United States Department of Agriculture, Professor Edward L. Greene of the United States National Museum, Professor Byron D. Halsted of Rutgers College and Professor William Trelease of the Missouri Botanical Garden have consented to act as an advisory committee.

Each author will be wholly responsible for his own contributions, being restricted only to the general style adopted for the work, which must vary somewhat in the treatment of diverse groups.

The subscription price is fixed at $1.50 for each part; it is expected that four or five parts will be required for each volume. A limited number of separate parts will be sold at $2.00 each. Address,

THE NEW YORK BOTANICAL GARDEN

BRONX PARK, NEW YORK CITY

Volume 7, Part 1, Ustilaginales, including Ustilaginaceae and Tilletiaceae, by G. P. Clinton, was issued Oct. 4, 1906.

Volume 7, Part 2, commencing the Uredinales, by J. C. Arthur, is in press

Volume 22, Part 1, including Podostemonaceae by George V. Nash, Crassulaceae by N. L. Britton and J. N. Rose, and Penthoraceae and Parnassiaceae by P. A. Rydberg, was issued May 22, 1905.

Volume 22, Part 2, including Saxifragaceae and Hydrangeaceae by J. K. Small and P. A. Rydberg, Cunoniaceae, Iteaceae and Hamamelidaceae by N. L. Britton, Pterostemonaceae by J. K. Small, Altingiaceae by Percy Wilson and Phyllonomaceae by H. H. Rusby, was issued Dec. 18, 1905.

PUBLICATIONS BY OFFICERS AND STUDENTS OF THE DEPARTMENT OF BOTANY, COLUMBIA UNIVERSITY

1. MEMOIRS OF THE DEPARTMENT:

Vol. 1. A Monograph of the North American Species of the Genus *Polygonum* (1895). By John Kunkel Small, Fellow in Botany, 1893–1895; Curator of the Herbarium, 1895–1898.

Quarto, 178 pages, 84 plates. Price $6.00.

Vol. 2. A Monograph of the North American Potentilleae (1898). By Per Axel Rydberg, Fellow in Botany, 1896–7.

Quarto, 224 pages, 112 plates. Price $6.00.

2. CONTRIBUTIONS FROM THE DEPARTMENT:

Vol. 1. Nos. 1–25. 1886–1892. Price $5.00.
Vol. 2. Nos. 26–50. 1892–1894. Price $5.00.
Vol. 3. Nos. 51–75. 1894–1895. Price $5.00.
Vol. 4. Nos. 76–100. 1895–1896. Price $5.00.
Vol. 5. Nos. 101–125. 1896–1897. Price $5.00.
Vol. 6. Nos. 126–150. 1897–1898. Price $5.00.
Vol. 7. Nos. 151–175. 1898–1901. Price $5.00.
Vol. 8. Nos. 176–200. 1901–1902. Price $5.00.
Vol. 9. Nos. 201–. 1903–(current).

List of separate numbers available on application.

3. A Text-book of General Lichenology (1896). By Albert Schneider, Fellow in Botany, 1895–1896.

Octavo, 230 pages, 76 plates. Price $4.25 (cloth); $3.80 (paper). Published by Willard N. Clute & Co. Binghamton, N. Y.

The Department invites propositions relative to exchanges of herbarium material for any of the above named publications. Address Professor L. M. Underwood, Columbia University, N. Y. City.

4. An Illustrated Flora of the Northern United States, Canada, and the British Possessions from Newfoundland to the Parallel of the Southern Boundary of Virginia, and from the Atlantic Ocean westward to the 102d Meridian (1896–1898). By Nathaniel Lord Britton, Professor of Botany, 1891–1896; Emeritus Professor 1896–, and Hon. Addison Brown.

Three volumes, royal octavo. Vol. 1, 612 pages; Vol. 2, 642 pages; Vol. 3, 588 pages; 4162 figures in the text, illustrating every species described.

Published by Charles Scribner's Sons, New York. Price, in cloth, $9.00 for the three volumes; with the indexes and keys bound separately, $10.00.

5. Our Native Ferns and their Allies. (Sixth Edition) 1900. By Lucien Marcus Underwood, Professor of Botany, 1896–.

Duodecimo, 158 pages, 35 figures. Published by Henry Holt & Co., New York. Price $1.00.

6. Moulds, Mildews, and Mushrooms. By Lucien Marcus Underwood, Professor of Botany, 1896—.

Duodecimo, 236 pages, 10 plates. Published by Henry Holt & Co. Price $1.50.

7. A Text-book of General Botany (1897). By Carlton Clarence Curtis, Assistant in Botany, 1892-1895; Tutor in Botany, 1895–.

Octavo, 360 pages; 87 illustrations. Published by Longmans, Green & Co., New York. Price $3.00.

ESTABLISHED 1851

EIMER & AMEND

205–211 Third Ave.,
Corner 18th Street NEW YORK

MANUFACTURERS AND IMPORTERS OF

Chemicals and Chemical Apparatus

Our Specialties

MICROSCOPES
and
ACCESSORIES

MICROTOMES

BALANCES
and
WEIGHTS

Sole Agents for
C. A. F.
KAHLBAUM'S
C. P. Chemicals
and Reagents

CARL
SCHLEICHER
& SCHÜLL'S
C. P. Filters

SCHOTT &
GENOSSEN'S
Famous Laboratory Glassware

And for many other renowned European houses in our line.

COMPLETE LINE OF

Bacteriological Apparatus

WE HANDLE

EVERYTHING NEEDED FOR A LABORATORY

Inspection of premises cordially invited.

MEMOIRS OF THE TORREY BOTANICAL CLUB

A series of technical papers on botanical subjects, published at irregular intervals. Price $3.00 a volume. Not offered in exchange.

Vol. 1, No. 1; price, $1.00:

Bailey, Liberty Hyde. Studies of the types of various species of the genus *Carex*. Pages 1–85. 25 My 1889.

Vol. 1, No. 2; not furnished separately:

Martindale, Isaac Comly. Marine algae of the New Jersey coast and adjacent waters of Staten Island. Pages 87–111. 24 Au 1889.

Vol. 1, No. 3; price, 75 cents:

Spruce, Richard. *Hepaticae* Bolivianae, in Andibus Boliviae orientalis, annis 1885–6, a cl. H. H. Rusby lectae. Pages 113–140. 20 Ja 1890.

Vol. 1, No. 4; price, 75 cents:

Sturtevant, Edward Lewis. On seedless fruits. Pages 141–187. 30 My 1890.

Vol. 2, No. 1; not furnished separately:

Halsted, Byron David. Reserve food-materials in buds and surrounding parts. Pages 1–26, *plates 1, 2*. 10 S 1890.

Vol. 2, No. 2; price, 75 cents:

Vail, Anna Murray & Hollick, Charles Arthur. Contributions to the botany of Virginia. Pages 27–56, *plates 3, 4*. 23 D 1890.

Vol. 2, No. 3; not furnished separately:

Holm, Herman Theodor. Contributions to the knowledge of the germination of some North American plants. Pages 57–108, *plates 5–19*. 15 Ap 1891.

Vol. 2, No. 4; price, 75 cents:

Wheelock, William Elmer. The genus *Polygala* in North America. Pages 109–132. 30 D 1891.

Vol. 3, No 1; not furnished separately:

Small, John Kunkel & Heller, Amos Arthur. Flora of western North Carolina and contiguous territory. Pages 1–39. 20 F 1892.

Vol 3, No. 2; price, $2.00:

Morong, Thomas. The *Naiadaceae* of North America [with illustrations of all the species]. Pages 1–65, *plates 20–74*. 15 Mr 1893.

Vol. 3, No. 3; not furnished separately:

Rusby, Henry Hurd. An enumeration of the plants collected in Bolivia by Miguel Bang. [Part 1.] Pages 1–67. 28 Ap 1893.

Vol. 4, No. 1; not furnished separately:

Underwood, Lucien Marcus. Index *Hepaticarum*. Part I, Bibliography. Pages 1–91. 10 Je 1893.

Vol. 4, No. 2; not furnished separately:

Small, John Kunkel & Vail, Anna Murray. Report on the botanical exploration of southwestern Virginia during the season of 1892. Pages 92–201, *plates 75–82*. 18 N 1893–17 Ap 1894.

Vol. 4, No. 3; price, 50 cents:

Rusby, Henry Hurd. An enumeration of the plants collected in Bolivia by Miguel Bang. Part 2. Pages 203–274. 10 Mr–17 Ap 1895.

Vol. 4, No. 4; price, 50 cents:

Pettit, Anna Stockton. *Arachis hypogaea* L. Pages 275–296, *plates 83–85*. 5 Je 1895.

Vol. 4, No. 5; price, 75 cents:

Rydberg, Per Axel. The North American species of *Physalis* and related genera. Pages 297–374. 15 S 1896.

Vol. 5; price, $3.00:

List of *Pteridophyta* and *Spermatophyta* growing without cultivation in northeastern North America. Pages 1–377. 4 D 1893–31 D 1894.

Vol. 6, No. 1; price, $1.25:

Rusby, Henry Hurd. An enumeration of the plants collected in Bolivia by Miguel Bang. Part 3. Pages 1–130. 17 N 1896.

Vol. 6, No. 2; price, 50 cents:

Grout, Abel Joel. A revision of the North American *Isotheciaceae* and *Brachythecia*. Pages 131–210. 30 Jl 1897.

Vol. 6, No. 3; price, 50 cents:

Hazen, Tracy Elliot. The life history of *Sphaerella lacustris* (*Haematococcus pluvialis*). Pages 211–246, *plates 86, 87* (*colored*). 8 Je 1899.

Vol. 6, No. 4; price, 50 cents:

Underwood, Lucien Marcus. A review of the genera of ferns proposed prior to 1832. Pages 247–283. 1 D 1899.

Vol. 6, No. 5; price, 25 cents:

Fink, Bruce. Notes on the lichen distribution in the upper Mississippi Valley. Pages 285–307. 1 D 1899.

Vol. 7; price, $3.00:

Howe, Marshall Avery. The *Hepaticae* and *Anthocerotes* of California. Pages 1–208, *plates 88–122*. 5 Au 1899.

Vol. 8, No. 1; price, $1.75:

Lloyd, Francis Ernest. The comparative embryology of the *Rubiaceae*. Part I. Pages 1–26, *plates 1–4*. 26 Au 1899. Part II. Pages 27–112, *plates 5–15*. 15 F 1902.

Vol. 8, No. 2; price, $1.00:

Evans, Alexander William. The *Lejeuneae* of the United States and Canada. Pages 113–183, *plates 16–22*. 15 F 1902.

Vol. 8, No. 3; price, 75 cents:

Britton, Elizabeth Gertrude & Taylor, Alexandrina. The life history of *Vittaria lineata*. Pages 185–211, *plates 23–31*. 30 Au 1902.

Vol. 9; price, $3.00:

Salmon, Ernest Stanley. A monograph of the *Erysiphaceae*. Pages 1–292, *plates 1–9*. 4 O 1900.

Vol. 10; price, $3.00:

Burgess, Edward Sandford. History of Pre-Clusian botany in its relation to *Aster*. Pages i–xii, 1–447. 22 N 1902.

Vol. 11, No. 1; price, $1.75:

Griffiths, David. The North American *Sordariaceae*. Pages 1–134, *plates 1–19*. 30 My 1901.

Vol. 11, No. 2; price, $1.75:

Hazen, Tracy Elliot. The *Ulothricaceae* and *Chaetophoraceae* of the United States. Pages 135–250, *plates 20–42*. 20 O 1902.

Vol. 12, No. 1; price, $1.50:

Northrop, Alice Rich. Flora of New Providence and Andros (Bahama Islands). Pages 1–98, *plates 1–19*. 10 D 1902.

Vol. 12, No. 2; price, $1.00:

Banker, Howard James. A contribution to a revision of the North American Hydnaceae. Pages 99–194. 13 Je 1906.

Vol. 13; price, $3.00:

Burgess, Edward Sandford. Species and variations of Biotian Asters, with discussion of variability in *Aster*. Pages i–xv, 1–419, *plates 1–13*. 15 Mr 1906.

Address: **TORREY BOTANICAL CLUB,**
Columbia University, New York City.